

LECTURE 2 Aug 24<sup>th</sup>, 2023

PART I Fundamental Concepts in Quantum Information

This Lecture Understanding and Measuring one qubit

- Qubits and QM Law 1 (Superposition)
- Measurements and QM Law 2 (Born's Rule)
  - └ standard
  - └ in a different basis
- Uncertainty Principle
- Global vs Relative Phase

RECALL Qubit quantum state in superposition of two basic states " $|0\rangle$ " and " $|1\rangle$ "

" $\alpha$  amplitude on  $|0\rangle$ ,  $\beta$  amplitude on  $|1\rangle$ "  
where  $\alpha, \beta$  are complex numbers satisfying  $|\alpha|^2 + |\beta|^2 = 1$   $\begin{pmatrix} \alpha \\ \beta \end{pmatrix} \in \mathbb{C}^2$

E.g. a photon may have the state " $\frac{1}{\sqrt{2}}$  amplitude on  $|0\rangle$ ,  $\frac{1}{\sqrt{2}}$  amplitude on  $|1\rangle$ "

OR


Is this a quantum state?  $\leftarrow$  " $\frac{i}{\sqrt{2}}$  amplitude on  $|0\rangle$ , 0 amplitude on  $|1\rangle$ "

OR

"1 amplitude on  $|0\rangle$ , 0 amplitude on  $|1\rangle$ "  
called " $|0\rangle$ "  
reverse " $|1\rangle$ "

You cannot read a quantum state, i.e., access  $\alpha, \beta$  directly  
Only way to extract information is via measurement

What happens if a photon in a superposition state goes into the measuring device?



$\alpha$  ampl. " $|0\rangle$ "  
 $\beta$  ampl. " $|1\rangle$ "

This scenario is described by the second law of quantum mechanics

QM Law 2  
(Born's Rule)


For a particle with  $\alpha$  amplitude on  $|0\rangle$ ,  $\beta$  amplitude on  $|1\rangle$   
if you measure it, then

w/prob  $|\alpha|^2$ , readout shows " $|0\rangle$ " AND state becomes "1 amplitude on  $|0\rangle$ "  
w/prob  $|\beta|^2$ , readout shows " $|1\rangle$ " state becomes "1 amplitude on  $|1\rangle$ "  
whatever outcome was observed


E.g. photon in state " $\frac{4}{5}$  amplitude on  $|0\rangle$ ,  $\frac{3}{5}$  amplitude on  $|1\rangle$ "

w.p. 0.64, readout shows  $|0\rangle$   
w.p. 0.36, readout shows  $|1\rangle$

Example



Example (Filter)



To describe Born's rule more generally, we first take a detour and introduce quantum notation

Recall


a qubit in state " $\alpha$  ampl. on  $|0\rangle$ ",  $\beta$  ampl. on  $|1\rangle$ "  
can be described as a column vector

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} \in \mathbb{C}^2 \text{ s.t. } |\alpha|^2 + |\beta|^2 = 1$$

It is also a unit vector

If we only use real amplitudes  $\alpha, \beta$

length of  $(\alpha, \beta)$   
 $= \sqrt{\alpha^2 + \beta^2} = 1$



$$|0\rangle = 1 \text{ ampl. on } |0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$|1\rangle = \text{reverse} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$\{|0\rangle, |1\rangle\}$  forms an orthonormal basis for  $\mathbb{R}^2$


Any qubit can be written as  $\alpha|0\rangle + \beta|1\rangle$  where  $|\alpha|^2 + |\beta|^2 = 1$

Let 
$$\alpha \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \text{"}\psi\text{"} = \text{unit vector}$$

How can we describe  $\alpha, \beta$  in terms of  $\psi$ ?

$$\alpha = \text{"Projection of } \psi \text{ on } |0\rangle\text{"} = \langle \psi, |0\rangle \rangle$$

$$\beta = \text{"Projection of } \psi \text{ on } |1\rangle\text{"} = \langle \psi, |1\rangle \rangle$$



If  $u, v \in \mathbb{R}^n$ , then

$$\langle u, v \rangle = \text{Inner product of } u \text{ and } v = \sum_{i=1}^n u_i v_i = (u_1 \dots u_n) \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} = u^T v$$

$$\|u\| = \text{length of } u = \sqrt{\langle u, u \rangle} = \sqrt{\sum_{i=1}^n u_i^2} \quad \leftarrow \text{if } \psi = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

$$\|\psi\| = \sqrt{\alpha^2 + \beta^2}$$

What happens with complex vectors  $u, v \in \mathbb{C}^n$ ?

$$\langle u, v \rangle = \sum_{i=1}^n \bar{u}_i v_i = (\bar{u}_1 \dots \bar{u}_n) \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} = u^\dagger v \quad \leftarrow \text{dagger or conjugate transpose of } u$$

$$\|u\| = \text{length} = \sqrt{\langle u, u \rangle} = \sqrt{\sum_{i=1}^n \bar{u}_i u_i} = \sqrt{\sum |u_i|^2}$$

So, if  $\psi = \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \in \mathbb{C}^2$ , then  $\|\psi\| = \sqrt{|\alpha|^2 + |\beta|^2}$

Also,  $\{|0\rangle, |1\rangle\}$  is an orthonormal basis for  $\mathbb{C}^2$

so,  $\alpha = \langle \psi, |0\rangle \rangle$  and  $\beta = \langle \psi, |1\rangle \rangle$

## Dirac's Bra-Ket Notation

- $|\psi\rangle =$  column vector e.g.  $|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

$|\cdot\rangle$  is called a ket

- $\langle\psi| =$  conjugate transpose of  $\psi = (\bar{\psi}_1 \dots \bar{\psi}_n)$   
 $\psi^\dagger$  row vector

$\langle\cdot|$  is called a bra

- Inner Product of  $|\psi\rangle, |\phi\rangle$

$$= (\bar{\psi}_1 \dots \bar{\psi}_n) \begin{pmatrix} \phi_1 \\ \vdots \\ \phi_n \end{pmatrix} = \langle\psi|\phi\rangle = \langle\psi|\phi\rangle$$

Given a qubit  $|\psi\rangle$  and orthonormal basis  $\{|0\rangle, |1\rangle\}$

$$|\psi\rangle = \underbrace{\langle 0|\psi\rangle}_{\text{scalar}} |0\rangle + \underbrace{\langle 1|\psi\rangle}_{\text{scalar}} |1\rangle$$



Can do the same for any basis  $\{|b_0\rangle, |b_1\rangle\}$

$$|\psi\rangle = \langle b_0|\psi\rangle |b_0\rangle + \langle b_1|\psi\rangle |b_1\rangle$$

Example

$$\begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle$$

= " $|+\rangle$ "



$$\begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{2}} |0\rangle - \frac{1}{\sqrt{2}} |1\rangle = \text{"}|-\rangle"$$

$$|0\rangle = \frac{1}{\sqrt{2}} |+\rangle + \frac{1}{\sqrt{2}} |-\rangle$$

$$|1\rangle = \frac{1}{\sqrt{2}} |+\rangle - \frac{1}{\sqrt{2}} |-\rangle$$

Exercise

How do you express  $\langle\psi|$  in terms of ket notation?

NEXT LECTURE : Born's rule for measurement in a different basis & more ...