


PART I Fundamental Concepts in Quantum Information



TODAY Quantum Circuits and Entanglement

RECAP Say Alice has a qubit $|\psi\rangle$ and Bob has a qubit $|\phi\rangle \in \mathbb{C}^2$

Question 1: What is the joint 4-d state?

Question 2: If Bob applies a unitary $U \in \mathbb{C}^{2 \times 2}$ to his qubit, what is the new 4-d state?

Question 3: If only Alice measures her qubit, what happens?

Joint state

$$\begin{array}{ccc}
 \text{Alice } d\text{-qudit} & & \text{Bob } e\text{-qudit} \\
 |\psi\rangle = \begin{array}{c} |1\rangle \\ \vdots \\ |d\rangle \end{array} \begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_d \end{bmatrix} & & |\phi\rangle = \begin{array}{c} |1\rangle \\ \vdots \\ |e\rangle \end{array} \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_e \end{bmatrix}
 \end{array}$$

Joint state is de -dimensional qudit

$$\begin{array}{c}
 |11\rangle \\
 |12\rangle \\
 \vdots \\
 |1e\rangle \\
 |21\rangle \\
 \vdots \\
 |de\rangle
 \end{array}
 \begin{bmatrix} \alpha_1 \beta_1 \\ \alpha_1 \beta_2 \\ \vdots \\ \alpha_1 \beta_e \\ \alpha_2 \beta_1 \\ \vdots \\ \alpha_d \beta_e \end{bmatrix}
 =
 \begin{bmatrix} \alpha_1 \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_e \end{bmatrix} \\ \alpha_2 \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_e \end{bmatrix} \\ \vdots \\ \alpha_d \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_e \end{bmatrix} \end{bmatrix}
 = |\psi\rangle \otimes |\phi\rangle$$

This operation is called a **tensor product**.

More generally, tensor product of two matrices A and B:

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix}$$

$m \times n$ matrix

$$B = \begin{bmatrix} b_{11} & \dots & b_{1q} \\ \vdots & & \vdots \\ b_{p1} & \dots & b_{pq} \end{bmatrix}$$

$p \times q$ matrix

$$A \otimes B = \begin{bmatrix} a_{11}B & a_{12}B & \dots & a_{1n}B \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}B & a_{m2}B & \dots & a_{mn}B \end{bmatrix}$$

Each block is a $p \times q$ matrix

$mp \times nq$ matrix

E.g. $|0\rangle \otimes |0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \begin{matrix} |00\rangle \\ |01\rangle \\ |10\rangle \\ |11\rangle \end{matrix} = |00\rangle$

$$|0\rangle \otimes |+\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} |00\rangle + \frac{1}{\sqrt{2}} |01\rangle$$

$\frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle$

$$|+\rangle \otimes |0\rangle = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} |00\rangle + \frac{1}{\sqrt{2}} |10\rangle$$

Demonstrates that tensor product is not a commutative operation

$$\bullet |+\rangle \otimes |-\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \otimes \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \end{bmatrix} \begin{matrix} |00\rangle \\ |01\rangle \\ |10\rangle \\ |11\rangle \end{matrix}$$

Let's do this in the ket notation

$$\left(\frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle \right) \otimes \left(\frac{1}{\sqrt{2}} |0\rangle - \frac{1}{\sqrt{2}} |1\rangle \right) = \frac{1}{2} |00\rangle - \frac{1}{2} |01\rangle + \frac{1}{2} |10\rangle - \frac{1}{2} |11\rangle$$

Properties of tensor product

- Acts like "non-commutative multiplication"

$$(A+B) \otimes C = A \otimes C + B \otimes C$$

$$A \otimes (B+C) = A \otimes B + A \otimes C$$

$$A \otimes (B \otimes C) = (A \otimes B) \otimes C = A \otimes B \otimes C$$

E.g.

Alice

$$\begin{bmatrix} \alpha_0 \\ \alpha_1 \end{bmatrix}$$

Bob

$$\begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix}$$

Charlie

$$\begin{bmatrix} \gamma_0 \\ \gamma_1 \end{bmatrix}$$

Joint state is

$$\begin{bmatrix} \alpha_0 \beta_0 \gamma_0 \\ \alpha_0 \beta_0 \gamma_1 \\ \vdots \\ \alpha_1 \beta_1 \gamma_1 \end{bmatrix} \begin{matrix} |000\rangle \\ |001\rangle \\ \vdots \\ |111\rangle \end{matrix}$$


$$(A \otimes B)^{\dagger} = A^{\dagger} \otimes B^{\dagger}$$

$$(A \otimes B) \cdot (C \otimes D) = (AC) \otimes (BD)$$

↑
matrix
multiplication

Quantum Circuits

Let's suppose Alice and Bob each prepared a qubit and got together

Alice 

These two particles have a joint state

Bob 

$$\alpha_{00} |00\rangle + \alpha_{01} |01\rangle + \alpha_{10} |10\rangle + \alpha_{11} |11\rangle$$


↑↑
Alice Bob

Let's imagine they go in a physical device that changes their state (jointly)

e.g. the device applies a CNOT operation



Definition (CNOT)



"If the control qubit is 0, do nothing
Else, apply a NOT to the target qubit"

Formally, $|00\rangle \rightarrow |00\rangle$ $|10\rangle \rightarrow |11\rangle$
 $|01\rangle \rightarrow |01\rangle$ $|11\rangle \rightarrow |10\rangle$


The matrix representation of CNOT is

$$\begin{matrix}
 & |00\rangle & |01\rangle & |10\rangle & |11\rangle \\
 |00\rangle & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} & \begin{bmatrix} \alpha_{00} \\ \alpha_{01} \\ \alpha_{10} \\ \alpha_{11} \end{bmatrix} & \longrightarrow & \begin{bmatrix} \alpha_{00} \\ \alpha_{01} \\ \alpha_{11} \\ \alpha_{10} \end{bmatrix} \\
 |01\rangle & & & & \\
 |10\rangle & & & & \\
 |11\rangle & & & &
 \end{matrix}$$


This is a permutation matrix, so it is easy to see that this is a unitary transformation

What is the joint state of Alice and Bob's qubit after CNOT?

We draw this operation as a "quantum circuit" diagram



Let's draw a more interesting quantum circuit now



RECALL
 $H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
 $|0\rangle \rightarrow |+\rangle$
 $|1\rangle \rightarrow |-\rangle$

What are the states at locations ①, ② and ③?

State at location ① : $|00\rangle$

at location ② : Alice only applies a gate to her qubit $H|0\rangle = |+\rangle$
so, state is

$$|+\rangle \otimes |0\rangle = \frac{1}{\sqrt{2}} |00\rangle + \frac{1}{\sqrt{2}} |10\rangle$$

at location ③ : CNOT swaps the amplitude of $|10\rangle$ & $|11\rangle$
so, state is

$$\frac{1}{\sqrt{2}} |00\rangle + \frac{1}{\sqrt{2}} |11\rangle \rightarrow \text{Bell state} \\ \text{OR EPR pair}$$

Theorem

Bell State is not of the form $|\psi\rangle \otimes |\phi\rangle$ for any $|\psi\rangle, |\phi\rangle \in \mathbb{C}^2$

This means that such states can only arise when the particles interact

Proof

$$\text{Let } |\psi\rangle = \begin{bmatrix} \alpha_0 \\ \alpha_1 \end{bmatrix} \quad \text{and} \quad |\phi\rangle = \begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix}$$

$$\text{Then } |\psi\rangle \otimes |\phi\rangle = \begin{bmatrix} \alpha_0 \beta_0 \\ \alpha_0 \beta_1 \\ \alpha_1 \beta_0 \\ \alpha_1 \beta_1 \end{bmatrix}$$

Observe that the product of amplitudes on $|01\rangle$ & $|10\rangle$
= product of amplitudes on $|00\rangle$ & $|11\rangle$

for any tensor product state

This is not true for the Bell state ■

Definition


A state on multiple qubits is called **entangled across a bipartition** (of the qubits) if it cannot be written as $|\psi\rangle \otimes |\phi\rangle$ for any $|\psi\rangle \otimes |\phi\rangle$

It is not obvious just by looking at a state if its entangled

E.g. Is $\frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle + \frac{1}{2} |10\rangle + \frac{1}{2} |11\rangle$ entangled?

$$= |+\rangle \otimes |+\rangle$$

Suppose Bob applies a NOT gate to her qubit \rightarrow This is 2×2 unitary
 How can we make sense of this?



At location ④: $\frac{1}{\sqrt{2}} |01\rangle + \frac{1}{\sqrt{2}} |10\rangle$

Suppose Alice now applies a H gate

At location ⑤: with amplitude $\frac{1}{\sqrt{2}}$, state is $|01\rangle$
Alice Bob
↑ ↑

$$\begin{aligned}
 \text{applying H to Alice's qubit gives } & (H|0\rangle) \otimes |1\rangle \\
 & = |+\rangle \otimes |1\rangle \\
 & = \frac{1}{\sqrt{2}} |0\rangle \otimes |1\rangle + \frac{1}{\sqrt{2}} |1\rangle \otimes |1\rangle \\
 & = \frac{1}{\sqrt{2}} |01\rangle + \frac{1}{\sqrt{2}} |11\rangle
 \end{aligned}$$

with $\frac{1}{\sqrt{2}}$ amplitude, state is $|10\rangle$

$$\begin{aligned}
 \text{applying H to Alice's qubit gives } & (H|1\rangle) \otimes |0\rangle \\
 & = |-\rangle \otimes |0\rangle \\
 & = \frac{1}{\sqrt{2}} |00\rangle - \frac{1}{\sqrt{2}} |10\rangle
 \end{aligned}$$

$$\begin{aligned}
 \text{Final state, } & \frac{1}{\sqrt{2}} \left(\frac{1}{\sqrt{2}} |01\rangle + \frac{1}{\sqrt{2}} |11\rangle \right) + \frac{1}{\sqrt{2}} \left(\frac{1}{\sqrt{2}} |00\rangle - \frac{1}{\sqrt{2}} |10\rangle \right) \\
 & = \frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle - \frac{1}{2} |10\rangle + \frac{1}{2} |11\rangle
 \end{aligned}$$

In general, say we have a 2-qubit state $\begin{bmatrix} \alpha_{00} \\ \alpha_{01} \\ \alpha_{10} \\ \alpha_{11} \end{bmatrix} \in \mathbb{C}^4$ and a 2×2 unitary $U = \begin{bmatrix} p & r \\ q & s \end{bmatrix}$

What is the state after we apply U to 2nd qubit?

WLOG we only need to figure out what happens on the basis states $|00\rangle, |01\rangle, |10\rangle, |11\rangle$

$$|00\rangle = |0\rangle \otimes |0\rangle$$

$$\xrightarrow[2^{\text{nd}} \text{ qubit}]{U \text{ on}} |0\rangle \otimes U|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \otimes \begin{bmatrix} p \\ q \end{bmatrix} = \begin{bmatrix} p \\ q \\ 0 \\ 0 \end{bmatrix}$$

$$|01\rangle = |0\rangle \otimes |1\rangle \xrightarrow{U} |0\rangle \otimes U|1\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \otimes \begin{bmatrix} r \\ s \end{bmatrix} = \begin{bmatrix} r \\ s \\ 0 \\ 0 \end{bmatrix}$$

$$|10\rangle \xrightarrow{U} \begin{bmatrix} 0 \\ 0 \\ p \\ q \end{bmatrix}, \quad |11\rangle \xrightarrow{U} \begin{bmatrix} 0 \\ 0 \\ r \\ s \end{bmatrix}$$


Matrix for the overall transformation is

$$\begin{matrix} & 00 & 01 & 10 & 11 \\ \begin{bmatrix} p & r & 0 & 0 \\ q & s & 0 & 0 \\ 0 & 0 & p & r \\ 0 & 0 & q & s \end{bmatrix} & = & I_{2 \times 2} \otimes U \end{matrix}$$

If you did U to 1st qubit and nothing to 2nd : $U \otimes I_{2 \times 2}$

and V to 2nd : $U \otimes V$

E.g.



The order does not matter if you apply separate transformations on each particle