

### Homework #3

Please present 4 significant figures in your final answers for probabilities

1. (20 points) The joint probability mass function of discrete random variables X and Y taking values  $x = 1, 2, 3$  and  $y = 1, 2, 3$ , respectively, is given by a formula  $f_{XY}(x, y) = c \cdot (x + y)$ . Determine the following:

- a) (2 points) Find c

Answer:  $\sum_R f(x, y) = c \cdot (2 + 3 + 4 + 3 + 4 + 5 + 4 + 5 + 6) = 1$ ,  $c \cdot 36 = 1$ . Thus,  $c = 1/36$

- b) (2 points) Find probability of the event where  $X = 1$  and  $Y \leq 2$

Answer:  $P(X = 1, Y \leq 2) = f_{XY}(1,1) + f_{XY}(1,2) = \frac{1}{36}(2 + 3) = 5/36$

- c) (2 points) Find marginal probability  $P_Y(Y = 3)$

Answers:  $P(Y=3) = f_{XY}(1,3) + f_{XY}(2,3) + f_{XY}(3,3) = \frac{1}{36}(4 + 5 + 6) = 0.4167$

- d) (2 points) Marginal probability distribution of the random variable X

Answers: marginal distribution of X

x	$f_X(x) = f_{XY}(x,1) + f_{XY}(x,2) + f_{XY}(x,3)$
1	1/4
2	1/3
3	5/12

- e) (2 points)  $E(X)$ ,  $E(Y)$ ,  $V(X)$ , and  $V(Y)$

Answers:

$E(X) = \left(1 \times \frac{1}{4}\right) + \left(2 \times \frac{1}{3}\right) + \left(3 \times \frac{5}{12}\right) = 13/6 = 2.167$

$V(X) = E(X = 1) \cdot (1 - 2.167)^2 + E(X = 2) \cdot (2 - 2.167)^2 + E(X = 3) \cdot (3 - 2.167)^2 = 0.6389$

$E(Y) = 2.167$

$V(Y) = 0.6389$

- f) (2 points) Find conditional probability distribution of Y given that  $X = 1$

Answers:  $f_{Y|X}(y) = \frac{f_{XY}(1,y)}{f_X(1)}$

y	$f_{Y X}(y)$
1	$(2/36)/(1/4)=2/9$
2	$(3/36)/(1/4)=1/3$
3	$(4/36)/(1/4)=4/9$

- g) (2 points) Conditional probability distribution of X given that  $Y = 3$

Answers:  $f_{X|Y=3}(x) = \frac{f_{XY}(x,3)}{f_Y(3)}$  and  $f_Y(3) = f_{XY}(1,3) + f_{XY}(2,3) + f_{XY}(3,3) = \frac{1}{36}(4 + 5 + 6)$

$$= 0.4167$$

x	$f_{X Y}(x)$
1	$(4/36)/(15/36)=0.2667$
2	$(5/36)/(15/36)=0.3333$
3	$(6/36)/(15/36)=0.4$

h) (2 points) Are X and Y independent?

Answers: Since  $f_{XY}(1,1)=2/36 \neq 9/36 \cdot 9/36=2f_X(x)f_Y(y)$ , X and Y are not independent.

i) (2 points) What is the covariance for X and Y?

Answers:  $\text{cov}(X,Y) = \langle XY \rangle - \langle X \rangle \langle Y \rangle = (1/36) \cdot (2 \cdot 1 + 3 \cdot 2 + 4 \cdot 3 + 3 \cdot 2 + 4 \cdot 4 + 5 \cdot 6 + 4 \cdot 3 + 5 \cdot 6 + 6 \cdot 9) - 2.167 \cdot 2.167 = -0.0292$

j) (2 points) What is the correlation for X and Y?

Answers:  $\text{corr}(X,Y) = -0.0292/0.6389 = -0.0457$

2. (6 points) A random variable X has density function  $f(X = x) = c(x + x^3)$  for  $x \in [0,1]$  and  $f(X = x) = 0$  otherwise.

a) (3 points) Determine c.

Answer:  $c = 4/3$ .

b) (3 points) Compute  $E(1/X)$

Answer:  $E(1/X) = 16/9$

3. (10 points) Toss an unfair coin 4 times. The probability of head is 0.6 and that of tail is 0.4. Let X be the total number of heads among the first two tosses and Y the total number of heads among the last two tosses.

a) (4 points) Write down the joint probability mass fraction of X and Y.

Answers:

x/y	0	1	2	Margin
0	$0.4^4 = 0.0256$	$0.4^2 \times 2 \times 0.6 \times 0.4 = 0.0768$	$0.4^2 \times 0.6^2 = 0.0576$	0.16
1	$2 \times 0.6 \times 0.4 \times 0.4^2 = 0.0768$	$2 \times 0.4 \times 0.6 \times 2 \times 0.6 \times 0.4 = 0.2304$	$2 \times 0.6 \times 0.4 \times 0.6^2 = 0.1728$	0.48
2	$0.6^2 \times 0.4^2 = 0.0576$	$0.6^2 \times 2 \times 0.6 \times 0.4 = 0.1728$	$0.6^4 = 0.1296$	0.36
Margin	0.16	0.48	0.36	1.00

b) (2 points) Are X and Y independent? Please explain.

Answers: Independent.

- c) **(4 points)** Compute the conditional probability  $P(X \geq Y | X \geq 1)$

$$\text{Answers: } P(X \geq Y | X \geq 1) = \frac{P(X \geq Y, X \geq 1)}{P(X \geq 1)} = \frac{0.0768 + 0.2304 + 0.1296 + 0.1728 + 0.0576}{0.48 + 0.36} = 0.7943$$

4. **(6 points)** A random variable  $X$  is the average of  $p$  independent random variables  $X_k$ , i.e.,  $X = \frac{1}{p} \sum_{k=0}^p X_k$ , Calculate the expectation and the variance of  $X$  for three different cases:

- a) **(2 points)** When all  $X_k$  are independent uniform continuous random variables in the interval  $(0,1)$

$$\text{Answers: } E(X) = \frac{1}{2}, V(X) = \frac{1}{12p}$$

- b) **(2 points)** When all  $X_k$  are independent exponential random variables with PDF  $P(X_k = x) = \lambda_k e^{-\lambda_k x}$

$$\text{Answers: } E(X) = \frac{1}{p} \sum_k \lambda_k^{-1}, V(X) = \frac{1}{p^2} \sum_k \lambda_k^{-2}$$

- c) **(2 points)** When all  $X_k$  are Independent normal random variables but each one has its own mean  $\mu_k$  and its own standard deviation  $\sigma_k$

$$\text{Answers: } E(X) = \frac{1}{p} \sum_k \mu_k, V(X) = \frac{1}{p^2} \sum_k \sigma_k^2$$

5. **(4 points)** Suppose random variables  $X, Y$  have standard derivations,  $\sigma_X = 2$  and  $\sigma_Y = 6$ , respectively, and correlation coefficient  $\text{corr}(X, Y) = -1/3$ .

- (a) **(2 points)** Find  $\text{cov}(X, Y)$ .

$$\text{Answer: } \text{cov}(X, Y) = \text{Corr}(X, Y) * \sigma_X * \sigma_Y = -4$$

- (b) **(2 points)** Find  $\text{Var}(3X - 2Y)$ .

$$\text{Answers: } \text{Var}(4X - 2Y) = 9 * \text{Var}(X) + 4 * \text{Var}(Y) - 12 * \text{Cov}(X, Y) = 9 * 4 + 4 * 36 - 12 * (-4) = 228$$

6. **(12 points)** Time interval separating subsequent bus arrivals at a stop is an exponential random variable with mean 20 minutes. Steve and Andrew work at the same place and each will be late to work unless they board a bus on or before 8:40am. Steve comes to the bus stop exactly at 8am. Andrew also comes to the same bus stop but at a random time, uniformly distributed between 8am and 8:30am. Both of them take the first bus that arrives.

- (a) **(4 points)** What is the probability that Steve will be late for work tomorrow?

$$\text{Answers: } P(\text{Steve late}) = 1 - P(T < 40) = 1 - \frac{1}{20} \int_0^{40} e^{-t/20} dt = e^{-2} = 0.1353$$

- (b) **(4 points)** What is the probability that Andrew will be late for work tomorrow?

$$\text{Answers: } P(\text{Andrew late}) = \int_0^{30} \frac{dx}{30} P(T \geq 40 | T > x) = \int_0^{30} \frac{dx}{30} e^{-(40-x)/20} = \frac{e^{-2}}{30} \int_0^{30} e^{x/20} dx = \frac{20e^{-2}}{30} (e^{30/20} - 1) = 0.3141$$

(c) **(4 points)** What is the probability that Steve and Andrew will ride the same bus?

Probability that Steve will not leave by the time  $x$  when Andrew comes is  $\exp(-x/20)$ .

It needs to be integrated over  $\int_0^{30} \frac{dx}{30} \exp(-x/20) =$

$$\text{Answers: } P(\text{Steve and Andrew meet}) = \int_0^{30} \frac{dx}{30} e^{-x/20} = \frac{20}{30} (1 - e^{-30/20}) = 0.5179$$