Proving that a problem *X* is NP-hard requires several steps:

- Choose a problem *Y* that you already know is NP-hard (because we told you so in class).
- Describe an algorithm to solve *Y*, using an algorithm for *X* as a subroutine. Typically this algorithm has the following form: Given an instance of *Y*, transform it into an instance of *X*, and then call the magic black-box algorithm for *X*.
- *Prove* that your algorithm is correct. This always requires two separate steps, which are usually of the following form:
 - *Prove* that your algorithm transforms "good" instances of *Y* into "good" instances of *X*.
 - *Prove* that your algorithm transforms "bad" instances of *Y* into "bad" instances of *X*.
 Equivalently: Prove that if your transformation produces a "good" instance of *X*, then it was given a "good" instance of *Y*.
- Argue that your algorithm for *Y* runs in polynomial time. (This is usually trivial.)
- 1. Suppose you are given a magic black box that somehow answers the following decision problem in *polynomial time*:
 - INPUT: A boolean circuit *K* with *n* inputs and one output.
 - OUTPUT: TRUE if there are input values $x_1, x_2, ..., x_n \in \{\text{TRUE}, \text{FALSE}\}$ that make *K* output TRUE, and FALSE otherwise.

Using this black box as a subroutine, describe an algorithm that solves the following related search problem *in polynomial time*:

- INPUT: A boolean circuit *K* with *n* inputs and one output.
- OUTPUT: Input values $x_1, x_2, ..., x_n \in \{\text{TRUE}, \text{FALSE}\}$ that make *K* output TRUE, or NONE if there are no such inputs.


[Hint: You can use the magic box more than once.]

2. A *Hamiltonian cycle* in a graph *G* is a cycle that goes through every vertex of *G* exactly once. Deciding whether an arbitrary graph contains a Hamiltonian cycle is NP-hard.

A *tonian cycle* in a graph *G* is a cycle that goes through at least *half* of the vertices of *G*. Prove that deciding whether a graph contains a tonian cycle is NP-hard.

To think about later:

3. Let *G* be an undirected graph with weighted edges. A Hamiltonian cycle in *G* is *heavy* if the total weight of edges in the cycle is at least half of the total weight of all edges in *G*. Prove that deciding whether a graph contains a heavy Hamiltonian cycle is NP-hard.



A heavy Hamiltonian cycle. The cycle has total weight 34; the graph has total weight 67.