

Programming Languages and Compilers (CS 421)

Elsa L Gunter

2112 SC, UIUC

<http://courses.engr.illinois.edu/cs421>

Based in part on slides by Mattox Beckman, as updated by Vikram Adve and Gul Agha

3/5/24

1

Two Problems

Type checking

- Question: Does exp. e have type τ in env Γ ?
- Answer: Yes / No
- Method: Type derivation

Typability

- Question Does exp. e have some type in env. Γ ? If so, what is it?
- Answer: Type τ / error
- Method: Type inference

3/5/24

2

Type Inference - Outline

- Begin by assigning a type variable as the type of the whole expression
- Decompose the expression into component expressions
- Use typing rules to generate constraints on components and whole
- Recursively find substitution that solves typing judgment of first subcomponent
- Apply substitution to next subcomponent and find substitution solving it; compose with first, etc.
- Apply comp of all substitution to orig. type var. to get answer

3/5/24

3

Type Inference - Example

- What type can we give to $(\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x))$
- Start with a type variable and then look at the way the term is constructed

3/5/24

4

Type Inference - Example

- First approximate:
$$\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha$$
- Second approximate: use fun rule
$$\underline{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}$$

$$\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha$$
- Remember constraint $\alpha \equiv (\beta \rightarrow \gamma)$

3/5/24

5

Type Inference - Example

- Third approximate: use fun rule
$$\frac{\{f : \delta ; x : \beta\} \vdash f(f x) : \varepsilon}{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}$$

$$\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha$$
- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

6

Type Inference - Example

- Fourth approximate: use app rule

$$\frac{\{f:\delta; x:\beta\} \vdash f : \varphi \rightarrow \varepsilon \quad \{f:\delta; x:\beta\} \vdash f x : \varphi}{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

7

Type Inference - Example

- Fifth approximate: use var rule, get constraint $\delta \equiv \varphi \rightarrow \varepsilon$, Solve with same

- Apply to next sub-proof

$$\frac{\{f:\delta; x:\beta\} \vdash f : \varphi \rightarrow \varepsilon \quad \{f:\delta; x:\beta\} \vdash f x : \varphi}{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

8

Type Inference - Example

- Current subst: $\{\delta \equiv \varphi \rightarrow \varepsilon\}$

$$\frac{\dots \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f x : \varphi}{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

9

Type Inference - Example

- Current subst: $\{\delta \equiv \varphi \rightarrow \varepsilon\}$ Use App Rule

$$\frac{\{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f : \zeta \rightarrow \varphi \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash x : \zeta}{\dots \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f x : \varphi}$$

$$\frac{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

10

Type Inference - Example

- Current subst: $\{\delta \equiv \varphi \rightarrow \varepsilon\}$
- Var rule: Solve $\zeta \rightarrow \varphi \equiv \varphi \rightarrow \varepsilon$ Unification

$$\frac{\{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f : \zeta \rightarrow \varphi \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash x : \zeta}{\dots \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f x : \varphi}$$

$$\frac{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

11

Type Inference - Example

- Current subst: $\{\zeta \equiv \varepsilon, \varphi \equiv \varepsilon\} \circ \{\delta \equiv \varphi \rightarrow \varepsilon\}$
- Var rule: Solve $\zeta \rightarrow \varphi \equiv \varphi \rightarrow \varepsilon$ Unification

$$\frac{\{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f : \zeta \rightarrow \varphi \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash x : \zeta}{\dots \quad \{f:\varphi \rightarrow \varepsilon; x:\beta\} \vdash f x : \varphi}$$

$$\frac{\{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon}{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}$$

$$\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}$$

$$\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$$

3/5/24

12

Type Inference - Example

- Current subst: $\{\zeta \equiv \varepsilon, \varphi \equiv \varepsilon, \delta \equiv \varepsilon \rightarrow \varepsilon\}$

- Apply to next sub-proof

$$\begin{array}{c} \dots \quad \{f : \varepsilon \rightarrow \varepsilon; x : \beta\} \vdash x : \varepsilon \\ \hline \dots \quad \{f : \varphi \rightarrow \varepsilon; x : \beta\} \vdash f x : \varphi \\ \hline \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

13

Type Inference - Example

- Current subst: $\{\zeta \equiv \varepsilon, \varphi \equiv \varepsilon, \delta \equiv \varepsilon \rightarrow \varepsilon\}$

- Var rule: $\varepsilon \equiv \beta$

$$\begin{array}{c} \dots \quad \{f : \varepsilon \rightarrow \varepsilon; x : \beta\} \vdash x : \varepsilon \\ \hline \dots \quad \{f : \varphi \rightarrow \varepsilon; x : \beta\} \vdash f x : \varphi \\ \hline \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

14

Type Inference - Example

- Current subst: $\{\varepsilon \equiv \beta\} \circ \{\zeta \equiv \varepsilon, \varphi \equiv \varepsilon, \delta \equiv \varepsilon \rightarrow \varepsilon\}$

- Solves subproof; return one layer

$$\begin{array}{c} \dots \quad \{f : \varepsilon \rightarrow \varepsilon; x : \beta\} \vdash x : \varepsilon \\ \hline \dots \quad \{f : \varphi \rightarrow \varepsilon; x : \beta\} \vdash f x : \varphi \\ \hline \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

15

Type Inference - Example

- Current subst: $\{\varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$

- Solves this subproof; return one layer

$$\begin{array}{c} \dots \\ \dots \quad \{f : \varphi \rightarrow \varepsilon; x : \beta\} \vdash f x : \varphi \\ \hline \dots \quad \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \hline \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

16

Type Inference - Example

- Current subst: $\{\varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$

- Need to satisfy constraint $\gamma \equiv (\delta \rightarrow \varepsilon)$, given subst, becomes: $\gamma \equiv ((\beta \rightarrow \beta) \rightarrow \beta)$

$$\begin{array}{c} \dots \\ \hline \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

17

Type Inference - Example

- Current subst:

$$\{\gamma \equiv ((\beta \rightarrow \beta) \rightarrow \beta), \varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$$

- Solves subproof; return one layer

$$\begin{array}{c} \dots \\ \hline \{f : \delta ; x : \beta\} \vdash (f(f x)) : \varepsilon \\ \{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma \\ \{ \} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha \end{array}$$

- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

3/5/24

18

Type Inference - Example

- Current subst:

$$\{\gamma \equiv ((\beta \rightarrow \beta) \rightarrow \beta), \varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$$

- Need to satisfy constraint $\alpha \equiv (\beta \rightarrow \gamma)$
given subst: $\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \beta) \rightarrow \beta))$

$$\frac{\dots}{\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{\} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma);$

3/5/24

19

Type Inference - Example

- Current subst:

$$\{\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \beta) \rightarrow \beta)),$$

$$\gamma \equiv ((\beta \rightarrow \beta) \rightarrow \beta), \varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$$

- Solves subproof; return on layer

$$\frac{\dots}{\frac{\{x : \beta\} \vdash (\text{fun } f \rightarrow f(f x)) : \gamma}{\{\} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha}}$$

3/5/24

20

Type Inference - Example

- Current subst:

$$\{\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \beta) \rightarrow \beta)),$$

$$\gamma \equiv ((\beta \rightarrow \beta) \rightarrow \beta), \varepsilon \equiv \beta, \zeta \equiv \beta, \varphi \equiv \beta, \delta \equiv \beta \rightarrow \beta\}$$

- Done: $\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \beta) \rightarrow \beta))$

$$\{\} \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f(f x)) : \alpha$$

3/5/24

21

Type Inference Algorithm

Let $\text{infer}(\Gamma, e, \tau) = \sigma$

- Γ is a typing environment (giving polymorphic types to expression variables)
- e is an expression
- τ is a type (with type variables)
- σ is a substitution of types for type variables
- Idea: σ is substitution solving the constraints on type variables necessary for $\Gamma \vdash e : \tau$
- Should have $\sigma(\Gamma) \vdash e : \sigma(\tau)$ valid

3/5/24

22

Type Inference Algorithm

$\text{infer}(\Gamma, exp, \tau) =$

- Case exp of
 - **Var v** --> return $\text{Unify}\{\tau \equiv \text{freshInstance}(\Gamma(v))\}$
 - Replace all quantified type vars by fresh ones
 - **Const c** --> return $\text{Unify}\{\tau \equiv \text{freshInstance } \varphi\}$ where $\Gamma \vdash c : \varphi$ by the constant rules
 - **fun $x \rightarrow e$** -->
 - Let α, β be fresh variables
 - Let $\sigma = \text{infer}(\{x : \alpha\} + \Gamma, e, \beta)$
 - Return $\text{Unify}(\{\sigma(\tau) \equiv \sigma(\alpha \rightarrow \beta)\}) \circ \sigma$

3/5/24

23

Example of inference with Var Rule

Instance $\{a \rightarrow 'w\}$ ('w a fresh variable)

$\{x : \text{All } 'a. ('a * 'b) \text{ list}, y : \text{All. } 'b\} \vdash x : (\text{int} * \text{string}) \text{ list}$

$\text{freshInstance}(\text{All } 'a. ('a * 'b) \text{ list}) = ('w * 'b) \text{ list}$

$\text{Unify}\{((\text{int} * \text{string}) \text{ list}) = ('w * 'b) \text{ list}\} = \{'w \rightarrow \text{int}, 'b \rightarrow \text{string}\}$

After substitution:

Instance $\{a \rightarrow \text{int}\}$

$\{x : \text{All } 'a. ('a * \text{string}) \text{ list}, y : \text{All. string}\} \vdash x : (\text{int} * \text{string}) \text{ list}$

3/5/24

24

Type Inference Algorithm (cont)

- Case *exp* of
 - App ($e_1 e_2$) -->
 - Let α be a fresh variable
 - Let $\sigma_1 = \text{infer}(\Gamma, e_1, \alpha \rightarrow \tau)$
 - Let $\sigma_2 = \text{infer}(\sigma_1(\Gamma), e_2, \sigma_1(\alpha))$
 - Return $\sigma_2 \circ \sigma_1$

3/5/24

25

Type Inference Algorithm (cont)

- Case *exp* of
 - If e_1 then e_2 else e_3 -->
 - Let $\sigma_1 = \text{infer}(\Gamma, e_1, \text{bool})$
 - Let $\sigma_2 = \text{infer}(\sigma_1(\Gamma), e_2, \sigma_1(\tau))$
 - Let $\sigma_3 = \text{infer}(\sigma_2 \circ \sigma_1(\Gamma), e_3, \sigma_2 \circ \sigma_1(\tau))$
 - Return $\sigma_3 \circ \sigma_2 \circ \sigma_1$

3/5/24

26

Type Inference Algorithm (cont)

- Case *exp* of
 - let $x = e_1$ in e_2 -->
 - Let α be a fresh variable
 - Let $\sigma_1 = \text{infer}(\Gamma, e_1, \alpha)$
 - Let $\sigma_2 =$
 $\text{infer}(\{x: \text{GEN}(\sigma_1(\Gamma), \sigma_1(\alpha))\} + \sigma_1(\Gamma), e_2, \sigma_1(\tau))$
 - Return $\sigma_2 \circ \sigma_1$

3/5/24

27

Type Inference Algorithm (cont)

- Case *exp* of
 - let rec $x = e_1$ in e_2 -->
 - Let α be a fresh variable
 - Let $\sigma_1 = \text{infer}(\{x: \alpha\} + \Gamma, e_1, \alpha)$
 - Let $\sigma_2 = \text{infer}(\{x: \text{GEN}(\sigma_1(\Gamma), \sigma_1(\alpha))\} + \sigma_1(\Gamma), e_2, \sigma_1(\tau))$
 - Return $\sigma_2 \circ \sigma_1$

3/5/24

28

Type Inference Algorithm (cont)

- To infer a type, introduce *type_of*
- Let α be a fresh variable
- *type_of* (Γ, e) =
 - Let $\sigma = \text{infer}(\Gamma, e, \alpha)$
 - Return $\sigma(\alpha)$
- Need an algorithm for *Unif*

3/5/24

29

Background for Unification

- Terms made from constructors and variables (for the simple first order case)
- Constructors may be applied to arguments (other terms) to make new terms
- Variables and constructors with no arguments are base cases
- Constructors applied to different number of arguments (arity) considered different
- Substitution of terms for variables

3/5/24

30

Simple Implementation Background

```

type term = Variable of string
           | Const of (string * term list)
let x = Variable "a";; let tm = Const ("2",[]);;

let rec subst var_name residue term =
  match term with Variable name ->
    if var_name = name then residue else term
  | Const (c, tys) ->
    Const (c, List.map (subst var_name residue)
      tys);;
  
```

3/5/24

31

Unification Problem

Given a set of pairs of terms (“equations”)
 $\{(s_1, t_1), (s_2, t_2), \dots, (s_n, t_n)\}$
 (the *unification problem*) does there exist
 a substitution σ (the *unification solution*)
 of terms for variables such that
 $\sigma(s_i) = \sigma(t_i)$,
 for all $i = 1, \dots, n$?

3/5/24

32

Uses for Unification

- Type Inference and type checking
- Pattern matching as in OCaml
 - Can use a simplified version of algorithm
- Logic Programming - Prolog
- Simple parsing

3/5/24

33

Unification Algorithm

- Let $S = \{(s_1 = t_1), (s_2 = t_2), \dots, (s_n = t_n)\}$ be a unification problem.
- Case $S = \{\}$: $\text{Unif}(S) = \text{Identity function}$ (i.e., no substitution)
- Case $S = \{(s, t)\} \cup S'$: Four main steps

3/5/24

34

Unification Algorithm

- **Delete:** if $s = t$ (they are the same term) then $\text{Unif}(S) = \text{Unif}(S')$
- **Decompose:** if $s = f(q_1, \dots, q_m)$ and $t = f(r_1, \dots, r_m)$ (same f , same m !), then $\text{Unif}(S) = \text{Unif}(\{(q_1, r_1), \dots, (q_m, r_m)\} \cup S')$
- **Orient:** if $t = x$ is a variable, and s is not a variable, $\text{Unif}(S) = \text{Unif}(\{(x = s)\} \cup S')$

3/5/24

35

Unification Algorithm

- **Eliminate:** if $s = x$ is a variable, and x does not occur in t (the occurs check), then
 - Let $\varphi = \{x \rightarrow t\}$
 - $\text{Unif}(S) = \text{Unif}(\varphi(S')) \circ \{x \rightarrow t\}$
 - Let $\psi = \text{Unif}(\varphi(S'))$
 - $\text{Unif}(S) = \{x \rightarrow \psi(t)\} \circ \psi$
 - Note: $\{x \rightarrow a\} \circ \{y \rightarrow b\} = \{y \rightarrow (\{x \rightarrow a\}(b))\} \circ \{x \rightarrow a\}$ if y not in a

3/5/24

36

Tricks for Efficient Unification

- Don't return substitution, rather do it incrementally
- Make substitution be constant time
 - Requires implementation of terms to use mutable structures (or possibly lazy structures)
 - We won't discuss these

3/5/24

37

Example

- x, y, z variables, f, g constructors
- Unify $\{(f(x) = f(g(f(z), y))), (g(y, y) = x)\} = ?$

3/5/24

38

Example

- x, y, z variables, f, g constructors
- $S = \{(f(x) = f(g(f(z), y))), (g(y, y) = x)\}$ is nonempty
- Unify $\{(f(x) = f(g(f(z), y))), (g(y, y) = x)\} = ?$

3/5/24

39

Example

- x, y, z variables, f, g constructors
- Pick a pair: $(g(y, y) = x)$
- Unify $\{(f(x) = f(g(f(z), y))), (g(y, y) = x)\} = ?$

3/5/24

40

Example

- x, y, z variables, f, g constructors
- Pick a pair: $(g(y, y)) = x$
- Orient: $(x = g(y, y))$
- Unify $\{(f(x) = f(g(f(z), y))), (g(y, y) = x)\} = ?$
Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\}$
by Orient

3/5/24

41

Example

- x, y, z variables, f, g constructors
- Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\} = ?$

3/5/24

42

Example

- x, y, z variables, f, g constructors
- $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\}$ is non-empty
- Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\} = ?$

3/5/24

43

Example

- x, y, z variables, f, g constructors
- Pick a pair: $(x = g(y, y))$
- Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\} = ?$

3/5/24

44

Example

- x, y, z variables, f, g constructors
- Pick a pair: $(x = g(y, y))$
- Eliminate x with substitution $\{x \rightarrow g(y, y)\}$
 - Check: x not in $g(y, y)$
- Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\} = ?$

3/5/24

45

Example

- x, y, z variables, f, g constructors
- Pick a pair: $(x = g(y, y))$
- Eliminate x with substitution $\{x \rightarrow g(y, y)\}$
- Unify $\{(f(x) = f(g(f(z), y))), (x = g(y, y))\} =$
Unify $\{(f(g(y, y)) = f(g(f(z), y)))\}$
 - o $\{x \rightarrow g(y, y)\}$

3/5/24

46

Example

- x, y, z variables, f, g constructors
- Unify $\{(f(g(y, y)) = f(g(f(z), y)))\}$
 - o $\{x \rightarrow g(y, y)\} = ?$

3/5/24

47

Example

- x, y, z variables, f, g constructors
- $\{(f(g(y, y)) = f(g(f(z), y)))\}$ is non-empty
- Unify $\{(f(g(y, y)) = f(g(f(z), y)))\}$
 - o $\{x \rightarrow g(y, y)\} = ?$

3/5/24

48

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(f(g(y,y)) = f(g(f(z),y)))$
- Unify $\{(f(g(y,y)) = f(g(f(z),y)))\}$
 - o $\{x \rightarrow g(y,y)\} = ?$

3/5/24

49

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(f(g(y,y)) = f(g(f(z),y)))$
- Decompose: $(f(g(y,y)) = f(g(f(z),y)))$ becomes $\{(g(y,y) = g(f(z),y))\}$
- Unify $\{(f(g(y,y)) = f(g(f(z),y)))\}$
 - o $\{x \rightarrow g(y,y)\} =$
Unify $\{(g(y,y) = g(f(z),y))\} \circ \{x \rightarrow g(y,y)\}$

3/5/24

50

Example

- x,y,z variables, f,g constructors
- $\{(g(y,y) = g(f(z),y))\}$ is non-empty
- Unify $\{(g(y,y) = g(f(z),y))\}$
 - o $\{x \rightarrow g(y,y)\} = ?$

3/5/24

51

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(g(y,y) = g(f(z),y))$
- Unify $\{(g(y,y) = g(f(z),y))\}$
 - o $\{x \rightarrow g(y,y)\} = ?$

3/5/24

52

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(f(g(y,y)) = f(g(f(z),y)))$
- Decompose: $(g(y,y)) = g(f(z),y)$ becomes $\{(y = f(z)); (y = y)\}$
- Unify $\{(g(y,y) = g(f(z),y))\} \circ \{x \rightarrow g(y,y)\} =$
Unify $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\}$

3/5/24

53

Example

- x,y,z variables, f,g constructors
- Unify $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\} = ?$

3/5/24

54

Example

- x,y,z variables, f,g constructors
- $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\}$ is non-empty
- Unify $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\} = ?$

3/5/24

55

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(y = f(z))$
- Unify $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\} = ?$

3/5/24

56

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(y = f(z))$
- Eliminate y with $\{y \rightarrow f(z)\}$
- Unify $\{(y = f(z)); (y = y)\} \circ \{x \rightarrow g(y,y)\} =$
Unify $\{(f(z) = f(z))\}$
 - o $\{(y \rightarrow f(z)) \circ \{x \rightarrow g(y,y)\}\} =$
Unify $\{(f(z) = f(z))\}$
 - o $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\}$

3/5/24

57

Example

- x,y,z variables, f,g constructors
- Unify $\{(f(z) = f(z))\}$
 - o $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} = ?$

3/5/24

58

Example

- x,y,z variables, f,g constructors
- $\{(f(z) = f(z))\}$ is non-empty
- Unify $\{(f(z) = f(z))\}$
 - o $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} = ?$

3/5/24

59

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(f(z) = f(z))$
- Unify $\{(f(z) = f(z))\}$
 - o $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} = ?$

3/5/24

60

Example

- x,y,z variables, f,g constructors
- Pick a pair: $(f(z) = f(z))$
- Delete
- Unify $\{(f(z) = f(z))\}$
 - $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} =$
 - Unify $\{\} \circ \{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\}$

3/5/24

61

Example

- x,y,z variables, f,g constructors
- Unify $\{\} \circ \{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} = ?$

3/5/24

62

Example

- x,y,z variables, f,g constructors
- $\{\}$ is empty
- Unify $\{\} =$ identity function
- Unify $\{\} \circ \{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\} =$
 - $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\}$

3/5/24

63

Example

- Unify $\{(f(x) = f(g(f(z), y))), (g(y, y) = x)\} =$
 - $\{y \rightarrow f(z); x \rightarrow g(f(z), f(z))\}$
- $$\begin{aligned} f(\quad x \quad) &= f(g(f(z), \quad y \quad)) \\ \rightarrow f(g(f(z), f(z))) &= f(g(f(z), f(z))) \\ g(\quad y \quad , \quad y \quad) &= \quad x \\ \rightarrow g(f(z), f(z)) &= g(f(z), f(z)) \end{aligned}$$

3/5/24

64

Example of Failure: Decompose

- Unify $\{(f(x, g(y)) = f(h(y), x))\}$
- Decompose: $(f(x, g(y)) = f(h(y), x))$
- = Unify $\{(x = h(y)), (g(y) = x)\}$
- Orient: $(g(y) = x)$
- = Unify $\{(x = h(y)), (x = g(y))\}$
- Eliminate: $(x = h(y))$
- Unify $\{(h(y) = g(y))\} \circ \{x \rightarrow h(y)\}$
- No rule to apply! Decompose fails!

3/5/24

65

Example of Failure: Occurs Check

- Unify $\{(f(x, g(x)) = f(h(x), x))\}$
- Decompose: $(f(x, g(x)) = f(h(x), x))$
- = Unify $\{(x = h(x)), (g(x) = x)\}$
- Orient: $(g(x) = x)$
- = Unify $\{(x = h(x)), (x = g(x))\}$
- No rules apply.

3/5/24

66