

flag in advance; N is reset when such slips occur. The double-differenced pseudoranges are modelled in the same way, with the exception of the phase ambiguity term, which is not needed. Instead of solving for δx for all the stations independently, we solve for a single slip parameter $s(t)$, such that $\delta x(t) = Gs(t)$, where G are elastic functions relating slip on the model fault to displacement. We allow the slip to vary as a random walk in time, with scale σ_s . A nominal value of $\sigma_s = 3 \text{ mm day}^{-1/2}$ is small enough to limit the scatter during periods without anomalous deformation, but large enough to allow the full 87 mm of slip to accumulate over approximately 24 hours. The time-dependent model fits the GPS double-difference phases to within 4 mm (root mean square).

Received 30 August 2001; accepted 21 January 2002.

- McGuire, W. J. in *Volcano Instability on the Earth and Other Planets* (eds McGuire, W. J., Jones, A. P. & Neubeck, J.) 1–23 (Geological Society Spec. Publ. No. 110, London, 1996).
- Moore, J. G. et al. Prodigious submarine landslides on the Hawaiian Ridge. *J. Geophys. Res.* **94**, 17465–17484 (1989).
- Ward, S. Landslide tsunami. *J. Geophys. Res.* **106**, 11201–11216 (2001).
- Lipman, P. W., Normark, W. R., Moore, J. G., Wilson, J. B. & Gutmacher, C. E. The giant submarine Alike debris slide, Mauna Loa, Hawaii. *J. Geophys. Res.* **94**, 4279–4299 (1988).
- Cervelli, P., Murray, M. H., Segall, P., Aoki, Y. & Kato, T. Estimating source parameters from deformation data, with an application to the March 1997 earthquake swarm off the Izu Peninsula, Japan. *J. Geophys. Res.* **106**, 11217–11238 (2001).
- Efron, B. & Tibshirani, R. J. *An Introduction to the Bootstrap* (Chapman and Hall, New York, 1993).
- Linde, A. T., Gladwin, M. T., Johnston, M. J. S., Gwyther, R. L. & Bilham, R. G. A slow earthquake sequence on the San Andreas fault. *Nature* **383**, 65–68 (1996).
- Dragert, H., Wang, K. & James, T. S. A silent slip event on the deeper Cascadia subduction interface. *Science* **292**, 1525–1528 (2001).
- Nakamura, K. Why do long rift zones develop in Hawaiian volcanoes—A possible role of thick oceanic sediments. *Bull. Volcanol. Soc. Jpn* **25**, 255–267 (1980) (in Japanese).
- Dieterich, J. H. Growth and persistence of Hawaiian rift zones. *J. Geophys. Res.* **93**, 4258–4270 (1988).
- Thurber, C. H. & Gripp, A. E. Flexure and seismicity beneath the south flank of Kilauea Volcano and tectonic implications. *J. Geophys. Res.* **93**, 4271–4278 (1988).
- Delaney, P. T., Miklius, A., Árnadóttir, T., Okamura, A. T. & Sako, M. K. Motion of Kilauea Volcano during sustained eruption from the Pu'u O'o and Kupaianaha vents, 1983–1991. *J. Geophys. Res.* **98**, 17801–17820 (1993).
- Owen, S. et al. Rapid deformation of the south flank of Kilauea Volcano, Hawaii. *Science* **267**, 1328–1332 (1995).
- Owen, S. et al. Rapid deformation of Kilauea volcano: GPS measurements between 1990 and 1996. *J. Geophys. Res.* **105**, 18983–18998 (2000).
- Du, Y., Segall, P. & Gao, H. Quasi-static dislocations in three dimensional inhomogeneous media. *Geophys. Res. Lett.* **24**, 2347–2350 (1997).
- Aki, K. & Richards, P. G. *Quantitative Seismology; Theory and Methods* (Freeman, San Francisco, 1970).
- Ward, S. N. Quasi-static propagator matrices: creep on strike-slip faults. *Tectonophysics* **120**, 83–106 (1985).
- Okubo, P. G., Benz, H. M. & Chouet, B. A. Imaging the crustal magma sources beneath Mauna Loa and Kilauea volcanoes, Hawaii. *Geology* **25**, 867–870 (1997).
- Morgan, J. K., Moore, G. F., Hills, D. J. & Leslie, S. Overthrusting and sediment accretion along Kilauea's mobile south flank, Hawaii: Evidence for volcanic spreading from marine seismic reflection data. *Geology* **28**, 667–670 (2000).
- Elosegui, P., Davis, J., Johansson, J. & Shapiro, I. Detection of transient motions with the Global Positioning System. *J. Geophys. Res.* **101**, 11249–11261 (1996).
- Larson, K. et al. Volcano monitoring using kinematic GPS: Filtering strategies. *J. Geophys. Res.* **106**, 19453–19464 (2001).
- Stein, R. S. The role of stress transfer in earthquake occurrence. *Nature* **402**, 605–609 (1999).
- David, C., Wong, T.-F., Zhu, W. & Zhang, J. Laboratory measurements of compaction-induced permeability change in porous rocks; implications for the generation and maintenance of pore pressure excess in the crust. *Pure Appl. Geophys.* **143**, 425–456 (1994).
- Ingebritsen, S. E. & Scholl, M. A. The hydrology of Kilauea Volcano. *Geothermics* **22**, 255–270 (1993).
- Swanson, D. A., Duffield, W. A. & Fiske, R. S. Displacement of the south flank of Kilauea Volcano: The result of forceful intrusion of magma into the rift zones. *US Geol. Surv. Prof. Pap.* **963**, 1–30 (1976).
- Dvorak, J. J., Klein, F. W. & Swanson, D. A. Relaxation of the south flank after the 7.2-Magnitude Kalapana earthquake, Kilauea volcano, Hawaii. *Bull. Seismol. Soc. Am.* **84**, 133–141 (1994).
- Cannon, E. C., Bürgmann, R. & Owen, S. E. Shallow normal faulting and block rotation associated with the 1975 Kalapana earthquake, Kilauea Volcano, Hawaii. *Bull. Seismol. Soc. Am.* (in the press).
- Lichten, S. & Borden, J. Strategies for high-precision Global Positioning System orbit determination. *J. Geophys. Res.* **92**, 12751–12762 (1987).
- Hofmann-Wellenhof, B., Lichtenegger, H. & Collins, J. *GPS: Theory and Practice* (Springer, New York, 1994).

Acknowledgements

We thank M. Bevis for access to the GPS data for the University of Hawaii network, P. Okubo for providing hypocentres, G. Blewitt, R. Bürgmann, J. Davis, R. Iverson, M. Johnston, J. McGuire, J. Murray, D. Swanson and W. Thatcher for comments and discussion, and D. Okita for getting us to and from Kilauea GPS stations safely and efficiently. The National Science Foundation and the US Geological Survey provided funding for this research.

Competing interests statement

The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to P.C. (e-mail: cervelli@sumo.wr.usgs.gov).

Tyrannosaurus was not a fast runner

John R. Hutchinson*† & Mariano Garcia*†

* Department of Integrative Biology, University of California, Berkeley, California 94720-3140, USA

† Present addresses: Biomechanical Engineering Division, Stanford University, Stanford, California 94305-4038, USA (J.R.H.); Borg-Warner Automotive, 770 Warren Road, Ithaca, New York 14850, USA (M.G.)

The fastest gait and speed of the largest theropod (carnivorous) dinosaurs, such as *Tyrannosaurus*, is controversial. Some studies contend that *Tyrannosaurus* was limited to walking, or at best an 11 m s⁻¹ top speed^{1–4}, whereas others argue for at least 20 m s⁻¹ running speeds^{5–7}. We demonstrate a method of gauging running ability by estimating the minimum mass of extensor (supportive) muscle needed for fast running. The model's predictions are validated for living alligators and chickens. Applying the method to small dinosaurs corroborates other studies by showing that they could have been competent runners. However, models show that in order to run quickly, an adult *Tyrannosaurus* would have needed an unreasonably large mass of extensor muscle, even with generous assumptions. Therefore, it is doubtful that *Tyrannosaurus* and other huge dinosaurs (~6,000 kg) were capable runners or could reach high speeds.

Most assessments of running ability in large theropods are qualitative, based on analogies to walking elephants or running birds and hoofed mammals^{3,5–7}. Earlier quantitative assessments of the biomechanics of *Tyrannosaurus* suggested that it had limited locomotor performance^{1,2,4,8,9}, but uncertainties about tyrannosaur anatomy, physiology, and behaviour have impeded resolution of this debate^{1–9}. Fossilized footprints demonstrate that smaller theropod dinosaurs could run¹⁰, but running tracks from large theropods are unknown^{7,9}. Bulky limbs and other analogies with elephants




Figure 1 Explanation of free-body diagram analysis of body segments. Skeletal illustration modified from ref. 6. The initial *Tyrannosaurus* model (Trex_1) is shown in right lateral view and (x, y)-coordinate space, with the origin located at the toe joint. Pelvic pitch, hip, knee, ankle and toe joint angle definitions are shown (see Supplementary Information). One of the angles is redundant (four angles suffice). The ground reaction force (GRF) is vertical (typical for mid-stance^{12–14}) and acts at a distance R from the toe joints. The body segment weights (W_b , W_t , W_s and W_m for the trunk, thigh, shank and metatarsus) are also shown. A free-body diagram was used to calculate the internal moments about each joint. For example, about the knee joint (see inset), the moment that knee extensor muscles must generate (M_k) is equal to the ankle contact force ($-F_a$) times its respective moment arm, plus the gravitational moment of the shank segment and the ankle moment ($-M_a$).

provide evidence that the enormous sauropod dinosaurs did not run^{1,2,5,11}.

For extant animals, our results (Table 1) matched reality: alligators cannot run bipedally, whereas chickens are adept runners. Our dissected *Alligator* only had about 3.6% m_{body} per leg as extensor muscles, but we estimated that it would need about 7.7% m_{body} per leg to run quickly on its hindlimbs. In contrast, our *Gallus* specimen had about 8.8% m_{body} per leg as extensor muscles, but it should only need about 4.7% m_{body} per leg for fast running. Thus alligators have less than half the extensor muscle they would need to run fast, whereas chickens have almost twice the required amount of extensor muscle.

Well known scaling principles^{12–14} predict that animals of larger body mass (m_{body}) have more restricted locomotor performance because muscle force scales as $(m_{\text{body}})^{0.67}$, whereas muscle mass scales as $(m_{\text{body}})^{1.0}$. To be a fast runner, a *Tyrannosaurus* would have needed enough extensor muscle mass to support itself. We calculated (Fig. 1; Methods) this mass for various animals to determine whether running would require too much muscle mass.

Smaller theropods had lower estimates of minimum extensor muscle mass per leg (T) than *Tyrannosaurus* (Table 1), and thus we expect they had a broader range of running performance. We initially calculated (Fig. 2) that *Tyrannosaurus* needed about 43% of its body mass as extensor muscles in each leg (86% m_{body} for both

legs) in order to run quickly. This high estimate of T is surprising because our input parameter values were generously weighted towards a low T . These findings suggest that *Tyrannosaurus* could not run quickly.

A parameter study was then performed to assess the sensitivity of these findings to the values of the assumed parameters. Many of the parameters needed to model the limb mechanics of extinct dinosaurs are uncertain. To test how T was affected by the initial parameters, we entered a range of values and observed how T changed (Table 2). Some unknown parameters, such as the muscle fibre pennation angle (θ), were relatively unimportant: within a reasonable range of values (compared to extant taxa) they had little or no effect. Likewise, our approximations of the muscle moment arms (r) were presumably reliable because we used rigorous muscle reconstructions that minimize speculation^{15,16}. Although m_{body} is unknown for extinct taxa, our calculation expressed the required extensor muscle mass as a percentage of body mass (T), factoring out m_{body} (see Methods).

In contrast, uncertainty about other parameter values was quite important. The limb orientation used (Figs 1 and 2) was a critical assumption because a more columnar (straight-legged) orientation (Table 2) decreased the moment arms (R) of the ground reaction force (GRF) and hence decreased T compared to a more crouched (bent-legged) limb orientation. A more posterior position of the trunk centre of mass, shorter extensor muscle fibre lengths (L), greater extensor muscle moment arms (r), or a lower multiplier (G) of the GRF would also decrease our estimated values of T (Table 2). The relative lengths of muscle fibres (L) vary widely in extant taxa^{17,18}, so this parameter deserves more study. Admittedly, our calculations omitted other parameters that a more realistic model could include (Supplementary Information). However, most of these assumptions were conservative, leading to an underestimate of T .

We asked how much hindlimb muscle mass would be unreasonable for an extinct theropod dinosaur: in extant tetrapods, 50% or less of m_{body} is typically composed of muscle¹⁹. This proportion is

Table 1 Muscle anatomical data entered in equation (1) to estimate T

	Hip	Knee	Ankle	Toe
<i>Alligator</i>				
M_{muscle} (N m)	16	12	12	4.7
L (m)	0.092	–	0.037	0.0020
θ (°)	18	–	21	19
r (m)	0.019	–	0.017	0.0050
m_i (% m_{body})	4.9	0	1.7	1.1
T (% m_{body})				7.7
<i>Gallus</i>				
M_{muscle} (N m)	4.0	–0.30	4.7	2.2
L (m)	0.085	0.051	0.026	0.026
θ (°)	0	25	23	22
r (m)	0.038	0.026	0.010	0.0050
m_i (% m_{body})	0.08	2.0	2.0	
T (% m_{body})				4.7
<i>Coelophysis</i>				
M_{muscle} (N m)	45	–14	56	44
L (m)	0.17	0.097	0.065	0.037
θ (°)	9.0	21	22	21
r (m)	0.080	0.028	0.014	0.0060
m_i (% m_{body})	1.4	0.73	4.0	4.1
T (% m_{body})				10
Small tyrannosaur				
M_{muscle} (N m)	570	–180	580	390
L (m)	0.40	0.19	0.18	0.062
θ (°)	9.0	21	22	21
r (m)	0.14	0.057	0.025	0.020
m_i (% m_{body})	4.2	1.6	11.4	3.3
T (% m_{body})				21
<i>Tyrannosaurus</i> (Trex_1 best guess)				
M_{muscle} (N m)	75,000	–24,000	66,000	49,000
L (m)	1.2	0.52	0.39	0.19
θ (°)	9.0	21	22	21
r (m)	0.37	0.22	0.11	0.065
m_i (% m_{body})	15	3.6	15	9.0
T (% m_{body})				43

Values given are for all four limb joints (hip, knee, ankle, toe). M_{muscle} is the muscle moment required to maintain static equilibrium during single limb support at mid-stance of fast running (with $G = 2.5$); L is mean muscle fibre length; θ is mean extensor muscle fibre pennation angle; r is mean extensor muscle moment arm (means were weighted by physiological cross-sectional area as in refs 12–14); m_i is the percentage of m_{body} needed as extensor muscles about one joint; T is the total percentage of m_{body} per leg needed as extensor muscle (sum of all four m_i values).

The M_{muscle} shown is an extensor moment if it is positive, flexor if negative; except at the knee where the converse is true. In the *Alligator* model, the knee joint m_i was set at 0 because the knee M_{muscle} was a flexor moment and we conservatively assumed that two-joint hip extensors and knee flexors (for example *M. caudofemoralis longus*) could stabilize both joints. See Supplementary Information for more details.




Figure 2 Limb orientation used for model Trex_1. The values of m_i , the percentage of body mass as extensor muscle acting about each joint (muscle moment action shown as a red arrow) required to maintain static equilibrium at mid-stance during fast running are shown in red. Actual hindlimb bones were digitized from *Tyrannosaurus* specimen MOR 555 (Museum of the Rockies; Bozeman, Montana).

independent of m_{body} (ref. 20). Of that 50%, a total of 5–40% of m_{body} is allocated to limb extensor muscles^{15,17–19}. Considering these data, the simplest inference is that if T is greater than 25% m_{body} per leg as limb extensors (that is, more than 50% m_{body}), a biped could not run quickly (if at all), because it would not have enough muscle to exert the necessary forces.

Judging from extant taxa, values of T from 5–25% per leg indicate that an animal may or may not be a fast runner, depending on the ratio of the actual percentage of m_{body} that is extensor muscle to T , the estimated percentage of extensor muscle required. Values of T closer to 25% probably indicate poor running ability (if any) because the ratio of actual extensor muscle percentage to T would be perilously close to 1. Adept runners (such as ground birds^{12,21}) have T values below 5%, maintaining a high ratio of actual extensor muscle mass to T (almost 2.0 in *Gallus*). This allows the musculo-skeletal system to operate with a generous margin of safety^{12–14} to accommodate unexpected increases of joint moments.

Some robust conclusions are possible despite the uncertainties about parameter values. Our findings agree with data from fossilized footprints (fastest estimated speeds are $\sim 11 \text{ m s}^{-1}$)^{9,10} indicating that smaller theropods could run quickly. Although our T

estimates are somewhat high (10–21% m_{body} per leg), a more columnar limb orientation could easily have reduced T enough to enable fast running ($\sim 5\% m_{\text{body}}$).

Yet our estimates of T for an adult *Tyrannosaurus* were so high that it is difficult to justify the reconstruction of *Tyrannosaurus* as a fast runner. Even with generous assumptions we were unable to reduce the estimates of T to 5% m_{body} per leg or less. Indeed, scaling data^{17,18} suggest that a 6,000 kg *Tyrannosaurus* had only 7–10% m_{body} per leg as extensors. Therefore we conclude that an adult *Tyrannosaurus* had very limited, if any, running ability. Our best estimates indicate that an adult *Tyrannosaurus* needed over 40% m_{body} per leg as extensor muscle. If this is correct, then such animals were unable to run at all. If *Tyrannosaurus* was indeed an adept runner, then it must have had many musculoskeletal specializations that available data do not suggest.

A final model is illustrative. We isometrically scaled-up our chicken model to 6,000 kg to simulate a *Tyrannosaurus*-sized chicken. Our model shows that a gigantic chicken, using the same limb orientation as an extant galliform²¹, would need 99% m_{body} per leg as extensor muscles to run quickly, which is clearly impossible and is much higher than for a typical chicken (4.7% m_{body} per leg). Features such as the more posterior position of the trunk centre of mass for *Tyrannosaurus* explain why we obtained lower estimates of T for *Tyrannosaurus* (43% m_{body} per leg) than for the 6,000 kg chicken. Nonetheless, the high estimated values of T in both models reveal how giant terrestrial animals become restricted from extreme locomotor performance^{9,13}.

The ‘giant chicken’ example naturally prompts one to look at equation (1) in terms of scaling. Assuming isometric scaling, and that muscle pennation (θ) and stress (σ) are independent of body mass^{13,17,18,22}, we predict that T increases linearly with increasing size (Fig. 3). We expect T to increase with mass roughly in the manner shown by the dashed curve in Fig. 3, especially if limb orientation remains quite flexed.

Our conclusions lead us to question the reconstructions of *Tyrannosaurus* running at 11–20 m s^{-1} (refs 3–8). Even at a lower running speed of 11 m s^{-1} (refs 3, 4, 8), an adult *Tyrannosaurus* would have been near or above its maximum muscular capacity. A walking tyrannosaur could have adopted more extended limb joints

Table 2 Parameter sensitivity in *Tyrannosaurus* models

	Hip	Knee	Ankle	Toe
Trex_2				
M_{muscle} (N m)	53,000	–56,000	72,000	45,000
L (m)	1.2	0.52	0.39	0.19
θ (°)	9.0	21	22	21
r (m)	0.37	0.22	0.11	0.065
m_i (% m_{body})	10	8.3	16	8.3
T (% m_{body})				43
Trex_3				
M_{muscle} (N m)	75,000	49,000	80,000	29,000
L (m)	1.2	–	0.39	0.19
θ (°)	9.0	–	22	21
r (m)	0.37	–	0.11	0.065
m_i (% m_{body})	15	0	18	5.3
T (% m_{body})				38
Trex_4				
M_{muscle} (N m)	75,000	6,500	4,400	4,100
L (m)	1.2	–	0.39	0.19
θ (°)	9.0	–	22	21
r (m)	0.37	–	0.11	0.065
m_i (% m_{body})	15	0	0.99	0.75
T (% m_{body})				17
Trex_Lgator				
M_{muscle} (N m)	75,000	–24,000	66,000	49,000
L (m)	0.93	0.39	0.37	0.19
θ (°)	18	16	21	19
r (m)	0.37	0.22	0.11	0.065
m_i (% m_{body})	12	2.6	14	9.0
T (% m_{body})				38
Trex_Lchick				
M_{muscle} (N m)	75,000	–24,000	66,000	49,000
L (m)	1.1	0.65	0.41	0.33
θ (°)	0	25	23	22
r (m)	0.37	0.22	0.11	0.065
m_i (% m_{body})	13	4.6	16	16
T (% m_{body})				50
Trex_lowest				
M_{muscle} (N m)	75,000	6,500	4,400	4,100
L (m)	0.93	–	0.37	0.19
θ (°)	0	–	0	0
r (m)	0.37	–	0.11	0.065
m_i (% m_{body})	11	0	0.87	0.71
T (% m_{body})				13

This shows the effect on the estimate of T (relative to model Trex_1) when we changed limb orientation (models Trex_2–4) or muscle-fibre lengths (model Trex_Lgator and Trex_Lchick, isometrically scaling up L from *Alligator* (5.91 kg) and *Gallus* (2.89 kg) to *Tyrannosaurus* (6,000 kg). As with *Alligator* (Table 1), models Trex_3, Trex_4, and Trex_lowest had a flexor M_{muscle} about the knee joint, so the knee m_i was conservatively set at 0. Model Trex_lowest shows a combination of parameter values (L from model Trex_Lgator, limb orientation from model Trex_4, and θ set at 0°) that brought T to its lowest estimated value (see Supplementary Information).




Figure 3 Estimated extensor muscle mass (T , as a percentage of body mass per leg) required to run quickly. Data are from various models and extant species plotted against $\log(m_{\text{body}})$ to illustrate the scaling of T . Larger runners need a larger fraction of their body allocated to extensor muscle mass. Three shades of grey indicate the likelihood of running ability, based on estimated T . Very low estimated T implies a good chance of being a capable runner. Very high estimated T signals little chance of being a capable runner. Intermediate values of estimated T are inconclusive, and running ability in this case depends on the ratio of actual extensor muscle mass to T . Values of T greater than 50% per leg, or greater than 100% in total, are impossible. The solid line represents the case of scaling a chicken isometrically up to 6,000 kg.

to lower the GRF moment arms (R) and used double limb support to reduce the GRF, decreasing the T required for walking within reasonable bounds (Table 2: models *Trex_4* and *Trex_lowest*). However, our calculations suggest that even a walking tyrannosaur required activation of a large fraction of its extensor muscle volume, at considerable metabolic expense.

These conclusions should apply to all larger dinosaurs, invalidating arguments that tyrannosaurs were too slow to prey on large contemporaries³ such as *Triceratops*. *Tyrannosaurus* shows the most extreme cursorial (that is, locomotion-related) specializations of larger theropods^{5–9}, but even it was probably a slow runner, at best. Therefore we infer that the numerous lineages of theropod dinosaurs that evolved enormous sizes⁹ independently reduced their running ability. Conversely, reduction of body size and relative increases of extensor muscle sizes and moment arms probably increased running ability in maniraptoran theropods, especially birds^{15,16}. □

Methods

The minimum mass of muscle (m_i) required to produce the joint moment M_{muscle} to maintain static equilibrium about any joint i is a function of the muscle density (d), multiplied by the volume (V), divided by the fraction of active muscle volume (c). V is equal to the muscle fibre length (L), times the cross-sectional area (A), divided by the cosine of the pennation angle of the muscle fibres ($\cos\theta$)^{12–14}. Furthermore, A is equal to the muscle force (F) divided by the stress (σ). Finally, F must be the M_{muscle} required to stabilize each joint, divided by the mean extensor muscle moment arm (r)^{12–14}. Combining these expressions to solve for the extensor muscle mass required to act about a particular joint i (m_i ; expressed as a percentage of body mass, m_{body}) produces:

$$m_i = (100 M_{\text{muscle}} L d) / (\sigma c r m_{\text{body}} \cos\theta) \quad (1)$$

These parameters do not depend greatly on the physiology of the animal²², so the equation holds for both warm- and cold-blooded animals.

Estimating the joint moment M_{muscle}

In order to estimate M_{muscle} (and hence m_i), we used MATLAB (The Mathworks) software to compute the joint moments needed to maintain static equilibrium in a particular limb orientation. To maintain equilibrium, muscles must exert net muscle moments about joints that balance equal and opposite moments incurred by external and internal forces, such as the ground reaction force (GRF) and segment weights (W_{fb} , W_{c} , W_{s} and W_{m} in Fig. 1). These moments can be calculated in two dimensions by drawing a set of free-body diagrams and applying the three equilibrium equations ($\Sigma F_x = 0$, $\Sigma F_y = 0$, $\Sigma M = 0$). This procedure is appropriate for standing, and we explain below how it can be applied to running. We summed m_i for all four limb joints to estimate T , the total extensor muscle mass required per leg, as a percentage of m_{body} .

Choosing G

We multiplied the GRF magnitude in the free-body diagram analysis by a factor G , the ratio of the GRF to body weight ($m_{\text{body}}g$, with $g = 9.81 \text{ m s}^{-2}$), to estimate the M_{muscle} needed to support the body on one leg at mid-stance of fast running. At mid-stance of running, the body centre of mass is at its lowest point accelerating upwards, but inertial moments are minuscule relative to the moments of the GRF, so mid-stance is approximated as quasi-static^{12–14,23}. The value of G was set at 2.5, which is a reasonable value for fast running or hopping bipeds^{12–14,24,25}.

Determining forward velocity

To determine an approximate forward velocity that would be consistent with $G = 2.5$ for *Tyrannosaurus*, we can use the notion of dynamic similarity, such as the Froude number (Fr), the ratio of centripetal to gravitational forces²⁶, calculated as $Fr = (\text{velocity})^2 / (\text{hip height} \times g)$. Our value of $G = 2.5$ for *Tyrannosaurus* high-speed running is comparable to the value ($G = 2.7$) estimated for an ostrich²⁷ running at 12 m s^{-1} with $Fr = 16$. For *Tyrannosaurus* (hip height is roughly 2.5 m), $Fr = 16$ implies an average forward speed of almost 20 m s^{-1} . Therefore our model assumes a GRF magnitude consistent with the faster estimates of *Tyrannosaurus* running speed^{3–7}. In theory, at $Fr > 1$, an animal should switch from a walk to a run²⁶. Consequently the maximum walking speed that an adult tyrannosaur might have used was roughly 5 m s^{-1} (11 m.p.h.), which is still faster than speed estimates from known tracks of large theropods⁹.

Model parameters

To reconstruct running mechanics in extinct dinosaurs, we modelled a small adult theropod (*Coelophysis*), a small juvenile tyrannosaur, and an adult *Tyrannosaurus* (Supplementary Information). Functional segment lengths (distances between centres of joint rotation), masses, and centres of mass were measured from dissections for the extant taxa, or approximated for the extinct taxa. Hip, knee, ankle and toe joints connected these segments and were given joint angles. These parameters (see Supplementary Information) were used in a free-body diagram analysis (Fig. 1) to calculate M_{muscle} for equation (1).

We entered other parameter values (Table 1) into equation (1) to estimate T . The values of L , θ and r were measured from dissections for the extant taxa, or approximated for the extinct taxa. The muscle density (d) was set at $1.06 \times 10^3 \text{ kg m}^{-3}$ as in typical skeletal muscle²⁸, and the muscle stress (σ) was entered as $3.00 \times 10^7 \text{ N m}^{-2}$, the maximum isometric stress frequently measured²². Because we were estimating the minimum mass of muscle required to produce M_{muscle} , we set the fraction of active muscle volume (c) at 1. Muscle moment arms (r) were estimated for extinct taxa by applying muscle reconstructions^{15,16} to skeletal material, at appropriate joint angles.

T constraints

The ratio of actual extensor muscle mass to T , the required mass (both terms as a percentage of m_{body}) must be at least 1 for a bipedal animal to be able to run quickly; otherwise the actual muscles would be unable to generate the required forces. For extinct animals, we cannot measure the actual muscle masses, but as the estimated T increases, good running performance becomes increasingly unlikely. If T is more than 50% m_{body} per leg (100% m_{body} total), running at $Fr = 16$ is certainly impossible.

Received 27 July; accepted 12 December 2001.

- Alexander, R. McN. Mechanics of posture and gait of some large dinosaurs. *Zool. J. Linn. Soc.* **83**, 1–25 (1985).
- Alexander, R. McN. *Dynamics of Dinosaurs and Other Extinct Giants* (Columbia Univ. Press, New York, 1989).
- Hornor, J. R. & Lessem, D. *The Complete T. rex* (Simon & Schuster, New York, 1993).
- Farlow, J. O., Smith, M. B. & Robinson, J. M. Body mass, “strength indicator”, and cursorial potential of *Tyrannosaurus rex*. *J. Vert. Paleont.* **15**, 713–725 (1995).
- Bakker, R. T. *Dinosaur Heresies* (William Morrow, New York, 1986).
- Paul, G. S. *Predatory Dinosaurs of the World* (Simon & Schuster, New York, 1988).
- Paul, G. S. Limb design, function and running performance in ostrich-mimics and tyrannosaurs. *Gaia* **15**, 257–270 (2000).
- Christiansen, P. Strength indicator values of theropod long bones, with comments on limb proportions and cursorial potential. *Gaia* **15**, 241–255 (2000).
- Farlow, J. O., Gatesy, S. M., Holtz, T. R. Jr, Hutchinson, J. R. & Robinson, J. M. Theropod locomotion. *Am. Zool.* **40**, 640–663 (2000).
- Irby, G. V. Paleobiological evidence for running dinosaurs worldwide. *Mus. N. Arizona Bull.* **60**, 109–112 (1999).
- Coombs, W. P. Jr. Sauropod habits and habitats. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **17**, 1–33 (1975).
- Roberts, T. J. Muscle force and stress during running in dogs and wild turkeys. *Bull. Mus. Comp. Zool.* **156**, 283–295 (2001).
- Biewener, A. A. Scaling body support in mammals: limb posture and muscle mechanics. *Science* **245**, 45–48 (1989).
- Roberts, T. R., Chen, M. S. & Taylor, C. R. Energetics of bipedal running. II. Limb design and running mechanics. *J. Exp. Biol.* **201**, 2753–2762 (1998).
- Hutchinson, J. R. *The Evolution of Hindlimb Anatomy and Function in Theropod Dinosaurs*. PhD thesis, Univ. California (2001).
- Carrano, M. T. & Hutchinson, J. R. The pelvic and hind limb musculature of *Tyrannosaurus rex* (Dinosauria: Theropoda). *J. Morphol.* (in the press).
- Alexander, R. McN., Jayes, A. S., Maloij, G. M. O. & Wathuta, E. M. Allometry of the leg muscles of mammals. *J. Zool.* **194**, 539–552 (1981).
- Bennett, M. B. Allometry of the leg muscles of birds. *J. Zool.* **238**, 435–443 (1996).
- Grand, T. I. Body weight: its relation to tissue composition, segment distribution, and motor function. II. Interspecific comparisons. *Am. J. Phys. Anthropol.* **47**, 211–240 (1977).
- Hallgrímsson, B. & Maiorana, V. Variability and size in mammals and birds. *Biol. J. Linn. Soc.* **70**, 571–595 (2000).
- Gatesy, S. M. Guineafowl hind limb function. I. Cineradiographic analysis and speed effects. *J. Morphol.* **240**, 115–125 (1999).
- Johnston, I. A. Sustained force development: specializations and variation among the vertebrates. *J. Exp. Biol.* **115**, 239–251 (1985).
- Blickhan, R. The spring-mass model for running and hopping. *J. Biomech.* **22**, 1217–1227 (1989).
- Alexander, R. McN. The maximum forces exerted by animals. *J. Exp. Biol.* **115**, 231–238 (1985).
- Cavagna, G. A., Heglund, N. C. & Taylor, C. R. Mechanical work in terrestrial locomotion: two basic mechanisms for minimizing energy expenditure. *Am. J. Physiol.* **233**, 243–261 (1977).
- Alexander, R. McN. & Jayes, A. S. A dynamic similarity hypothesis for the gaits of quadrupedal mammals. *J. Zool.* **201**, 135–152 (1983).
- Alexander, R. McN., Maloij, G. M. O., Njau, R. & Jayes, A. S. Mechanics of running of the ostrich (*Struthio camelus*). *J. Zool.* **187**, 169–178 (1979).
- Mendez, J. & Keys, A. Density and composition of mammalian muscle. *Metabolism* **9**, 184–188 (1960).

Supplementary Information accompanies the paper on Nature’s website (<http://www.nature.com>).

Acknowledgements

We thank F. C. Anderson, K. Angielczyk, S. Delp, R. J. Full, S. M. Gatesy, T. M. Keaveny, R. Kram, K. Padian and J. F. Parham as well as the Berkeley Biomechanics and Stanford Neuromuscular Biomechanics seminars for input. J.R.H. appreciates funding from the University of California Department of Integrative Biology and Museum of Paleontology, and the National Science Foundation under a grant awarded in 2001. M.G. had postdoctorate funding through a DARPA grant to the Polypedal Lab at UC Berkeley. This is University of California Museum of Paleontology publication number 1749.

Correspondence and requests for materials should be addressed to J.R.H. (e-mail: jrhutch@stanford.edu).

6. Chan, F. K., Siegel, M. R. & Lenardo, J. M. Signaling by the TNF receptor superfamily and T cell homeostasis. *Immunity* **13**, 419–422 (2000).
7. Yang, Y., Fang, S., Jensen, J. P., Weissman, A. M. & Ashwell, J. D. Ubiquitin protein ligase activity of IAPs and their degradation in proteasomes in response to apoptotic stimuli. *Science* **288**, 874–877 (2000).
8. Tschopp, J., Martinon, F. & Hofmann, K. Apoptosis: Silencing the death receptors. *Curr. Biol.* **9**, R381–R384 (1999).
9. Van Antwerp, D. J., Martin, S. J., Kafri, T., Green, D. R. & Verma, I. M. Suppression of TNF- α -induced apoptosis by NF- κ B. *Science* **274**, 787–789 (1996).
10. Liu, Z. G., Hsu, H., Goeddel, D. V. & Karin, M. Dissection of TNF receptor 1 effector functions: JNK activation is not linked to apoptosis while NF- κ B activation prevents cell death. *Cell* **87**, 565–576 (1996).
11. Deveraux, Q. L. *et al.* IAPs block apoptotic events induced by caspase-8 and cytochrome *c* by direct inhibition of distinct caspases. *EMBO J.* **17**, 2215–2223 (1998).
12. Roy, N., Deveraux, Q. L., Takahashi, R., Salvesen, G. S. & Reed, J. C. The c-IAP-1 and c-IAP-2 proteins are direct inhibitors of specific caspases. *EMBO J.* **16**, 6914–6925 (1997).
13. Sarin, A., Conan-Cibotti, M. & Henkart, P. A. Cytotoxic effect of TNF and lymphotoxin on T lymphoblasts. *J. Immunol.* **155**, 3716–3718 (1995).
14. Zheng, L. *et al.* Induction of apoptosis in mature T cells by tumour necrosis factor. *Nature* **377**, 348–351 (1995).
15. Duckett, C. S. *et al.* A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors. *EMBO J.* **15**, 2685–2694 (1996).
16. Chan, F. K. & Lenardo, M. J. A crucial role for p80 TNF-R2 in amplifying p60 TNF-R1 apoptosis signals in T lymphocytes. *Eur. J. Immunol.* **30**, 652–660 (2000).
17. Arch, R. H., Gedrich, R. W. & Thompson, C. B. Translocation of TRAF proteins regulates apoptotic threshold of cells. *Biochem. Biophys. Res. Commun.* **272**, 936–945 (2000).
18. Huang, H. *et al.* The inhibitor of apoptosis, cIAP2, functions as a ubiquitin-protein ligase and promotes *in vitro* ubiquitination of caspases-3 and -7. *J. Biol. Chem.* **275**, 26661–26664 (2000).
19. Rothe, M., Wong, S. C., Henzel, W. J. & Goeddel, D. V. A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* **78**, 681–692 (1994).
20. Rothe, M., Sarma, V., Dixit, V. M. & Goeddel, D. V. TRAF2-mediated activation of NF- κ B by TNF receptor 2 and CD40. *Science* **269**, 1424–1427 (1995).
21. Reinhard, C., Shamoon, B., Shyamala, V. & Williams, L. T. Tumor necrosis factor α -induced activation of c-jun N-terminal kinase is mediated by TRAF2. *EMBO J.* **16**, 1080–1092 (1997).
22. Weiss, T. *et al.* Enhancement of TNF receptor p60-mediated cytotoxicity by TNF receptor p80: requirement of the TNF receptor-associated factor-2 binding site. *J. Immunol.* **158**, 2398–2404 (1997).
23. Weiss, T. *et al.* TNFR80-dependent enhancement of TNFR60-induced cell death is mediated by TNFR-associated factor 2 and is specific for TNFR60. *J. Immunol.* **161**, 3136–3142 (1998).
24. Erickson, S. L. *et al.* Decreased sensitivity to tumour-necrosis factor but normal T-cell development in TNF receptor-2-deficient mice. *Nature* **372**, 560–563 (1994).
25. Clem, R. J. *et al.* c-IAP1 is cleaved by caspases to produce a pro-apoptotic C-terminal fragment. *J. Biol. Chem.* **276**, 7602–7608 (2000).
26. Duckett, C. S. & Thompson, C. B. CD30-dependent degradation of TRAF2: implications for negative regulation of TRAF signaling and the control of cell survival. *Genes Dev.* **11**, 2810–2821 (1997).
27. Brown, K. D., Hostager, B. S. & Bishop, G. A. Differential signaling and tumor necrosis factor receptor-associated factor (traf) degradation mediated by CD40 and the Epstein-Barr virus oncoprotein latent membrane protein 1 (Lmp1). *J. Exp. Med.* **193**, 943–954 (2001).
28. Peschon, J. J. *et al.* TNF receptor-deficient mice reveal divergent roles for p55 and p75 in several models of inflammation. *J. Immunol.* **160**, 943–952 (1998).
29. Loric, K. L. *et al.* RING fingers mediate ubiquitin conjugating enzyme (E2)-dependent ubiquitination. *Proc. Natl Acad. Sci. USA* **96**, 11364–11369 (1999).
30. Memon, S. A., Petrak, D., Moreno, M. B. & Zacharchuk, C. M. A simple assay for examining the effect of transiently expressed genes on programmed cell death. *J. Immunol. Methods* **180**, 15–24 (1995).

Supplementary Information accompanies the paper on Nature's website (<http://www.nature.com>).

Acknowledgements

We are grateful to A. Weissman, C. Duckett, and Z. Liu for provision of reagents and reviews of this manuscript. X.L. is a visiting fellow from Bethune International Hospital, China.

Competing interests statement

The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to J.D.A. (e-mail: jda@pop.nci.nih.gov).

correction

Effects of experience and social context on prospective caching strategies by scrub jays

N. J. Emery & N. S. Clayton

Nature **414**, 443–446 (2001).

The exact number of species of American jays is a matter of debate¹. This is particularly evident within the scrub-jay lineage of *Aphelocoma*². Traditionally, *A. coerulescens* included the Florida scrub-jay and the western scrub-jay, among others. It has recently been brought to our attention that the Committee on Classification and Nomenclature of the American Ornithologists' Union now use *A. coerulescens* to refer exclusively to the Florida scrub-jay and *A. californica* to refer to the western scrub-jay². The birds used in the Emery & Clayton³ and Clayton & Dickinson⁴ studies were western scrub-jays: *A. californica*, not *A. coerulescens*. We therefore wish to correct the oversight in this paper and our other papers using these birds, and apologize for any confusion that has arisen. We thank R. Curry, T. Langen and G. Woolfenden for alerting us to the change in nomenclature. □

1. Madge, S. & Burn, H. *Crows and Jays. A Guide to the Crows, Jays and Magpies of the World* (Houghton Mifflin, New York, 1999).
2. American Ornithologists' Union *Check-list of North American Birds* 7th edn (American Ornithologists' Union, Washington DC, 1998).
3. Emery, N. J. & Clayton, N. S. Effects of experience and social context on prospective caching strategies by scrub jays. *Nature* **414**, 443–446 (2001).
4. Clayton, N. S. & Dickinson, A. Episodic-like memory during cache recovery by scrub jays. *Nature* **395**, 272–278 (1998).

erratum

Tyrannosaurus was not a fast runner

John R. Hutchinson & Mariano Garcia

Nature **415**, 1018–1021 (2002).

In this Letter, the row in Table 1 corresponding to 'Gallus, *m_i* (%*m_{body}*)' should read: 1.1, 0.08, 2.0 and 1.5 for the hip, knee, ankle and toe respectively. □