

$$|\vec{v}| \equiv \sqrt{\vec{v} \cdot \vec{v}} \quad \vec{v} = \sum_{i=1}^3 (\vec{v} \cdot \hat{r}_i) \hat{r}_i \quad d\vec{l}_{path} = \frac{d\vec{l}}{du} du \quad d\vec{A} = \left(\frac{\partial \vec{l}}{\partial u} \times \frac{\partial \vec{l}}{\partial v} \right) du dv$$

$$df(x_1, \dots, x_n) = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i \quad dV = \left(\frac{\partial \vec{l}}{\partial u} \times \frac{\partial \vec{l}}{\partial v} \right) \cdot \frac{\partial \vec{l}}{\partial w} du dv dw$$

$$\text{Conceptual version:} \quad d\vec{l}_{path} = d\vec{l}_u \quad d\vec{A} = d\vec{l}_u \times d\vec{l}_v$$

$$d\vec{l}_u \equiv \frac{\partial \vec{l}}{\partial u} du \quad dV = (d\vec{l}_u \times d\vec{l}_v) \cdot d\vec{l}_w$$

Taylor

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$


1st order approx for $x \ll 1$:

- $(1+x)^n \approx 1 + nx$
- $\sin x \approx x$
- $\cos x \approx 1 - \frac{x^2}{2}$
- $\tan x \approx x$
- $e^x \approx 1 + x$
- $\sin^{-1} x \approx x$
- $\cos^{-1} x \approx \frac{\pi}{2} - x$
- $\tan^{-1} x \approx x$
- $\ln(1+x) \approx x$

	0°	30°	45°	60°	90°
sin	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
tan	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞

Complex Numbers

$$e^{i\theta} = \cos \theta + i \sin \theta$$



$$\begin{aligned} \sin a \sin b &= \frac{1}{2} [\cos(a-b) - \cos(a+b)] \\ \cos a \cos b &= \frac{1}{2} [\cos(a+b) + \cos(a-b)] \\ \sin a \cos b &= \frac{1}{2} [\sin(a+b) + \sin(a-b)] \end{aligned}$$

$$\begin{aligned} \sin(a+b) &= \sin a \cos b + \cos a \sin b \\ \cos(a+b) &= \cos a \cos b - \sin a \sin b \end{aligned}$$

Integral Table

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \cos^{-1}\left(\frac{a}{x}\right) \quad \int \frac{dx}{\sqrt{x^2 \pm a^2}} = \ln\left(x + \sqrt{x^2 \pm a^2}\right) \quad \int \frac{x \, dx}{\sqrt{a^2 \pm x^2}} = \pm \sqrt{a^2 \pm x^2}$$

$$\int_0^{2\pi} \sin^2 \phi \, d\phi = \int_0^{2\pi} \cos^2 \phi \, d\phi = \pi \quad \int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1}\left(\frac{x}{a}\right) \quad \int \frac{x \, dx}{(a \pm x)^2} = \frac{a}{a \pm x} + \ln(a \pm x)$$

$$\int \sin^2 \phi \, d\phi = \frac{\phi}{2} - \frac{\sin(2\phi)}{4} \quad \int \frac{dx}{(a \pm x)^2} = \mp \frac{1}{a \pm x} \quad \int \frac{x \, dx}{a^2 \pm x^2} = \pm \frac{1}{2} \ln(a^2 \pm x^2)$$

$$\int \cos^2 \phi \, d\phi = \frac{\phi}{2} + \frac{\sin(2\phi)}{4} \quad \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) \quad \int \frac{x \, dx}{(a^2 \pm x^2)^{3/2}} = \mp \frac{1}{\sqrt{a^2 \pm x^2}}$$

$$\int \sin^3 \theta \, d\theta = \frac{\cos^3 \theta}{3} - \cos \theta \quad \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln\left(\frac{a+x}{a-x}\right) \quad \int \ln(ax) \, dx = x \ln(ax) - x$$

$$\int \cos^3 \theta \, d\theta = \sin \theta - \frac{\sin^3 \theta}{3} \quad \int \frac{dx}{(a^2 \pm x^2)^{3/2}} = \frac{x}{a^2 \sqrt{a^2 \pm x^2}} \quad \int \frac{\ln(ax)}{x} \, dx = \frac{1}{2} [\ln(ax)]^2$$

$$\int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{a^2 - x^2}}\right) \quad \int \frac{x^2}{\sqrt{a^2 - x^2}} \, dx = -\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{a^2 - x^2}}\right)$$

$$\int \sqrt{x^2 \pm a^2} \, dx = \frac{x}{2} \sqrt{x^2 \pm a^2} \pm \frac{a^2}{2} \ln|x + \sqrt{x^2 \pm a^2}| \quad \int \frac{(x - a \cos \theta) \sin \theta \, d\theta}{(x^2 + a^2 - 2ax \cos \theta)^{3/2}} = \frac{1}{x^2} \frac{a - x \cos \theta}{\sqrt{x^2 + a^2 - 2ax \cos \theta}}$$