Making Estimates in Research and Elsewhere Lance Cooper



assume rectangular barn w/ volume ~ LxW

 $N \sim (L \times W)/\pi R^2$

Making Estimates in Research: Why?

It is the mark of an instructed mind to rest assured with that degree of precision that the nature of the subject admits, and not to seek exactness when only an approximation of the truth is possible.

- Aristotle



Why make estimates in science?

The ability to estimate – to within an order of magnitude or so – the size or probability of various quantities is useful in science as well as in many other endeavors:

- To provide a rough check of more exact calculations
- To provide a rough check of research results or hypotheses
- To obtain estimates of quantities when other resources aren't available
- To obtain estimates of quantities that are difficult to measure precisely
- To obtain estimates of quantities for which no firm theoretical prediction exists ⇒ particularly important in interdisciplinary sciences, soft matter, astrophysics

To provide bounds for possible design alternatives

Making Estimates in Research: How?

How do you estimate the answer to a question that appears impossible to determine at all, or at least without access to an encyclopedia, internet connection, or omniscient being?

e.g., how many grains of sand are there on earth's beaches? how many piano tuners are there in Chicago? how many atoms are in your body?

These problems are sometimes referred to as Fermi problems, after the physicist Enrico Fermi, who was famous for (among other things) posing and solving such problems.

$-\frac{1}{U}\partial t = \frac{1}{2m} - \frac{1}{2}$ $\alpha = \frac{\pi^2}{ec}$

Enrico Fermi

Getting started:

- (1). Don't panic when you see the problem
- (2). Write down any fact you do know related to the question
- (3). Outline one or more possible procedures for determining the answer
- (4). List the things you'll need to know to answer the question
- (5). Keep track of your assumptions

Making Estimates in Research: How?

Other general guidelines for making order-of-magnitude estimates:

Make everything as simple as possible!

(1). Don't worry about specific values: round numbers to "convenient values"

e.g., $\pi \approx 3$; 8.4 ≈ 10 ; etc.

(2). Choose convenient geometries when modeling e.g., a spherical cow, a cubic grain of sand, etc.

(3). Make "educated" guesses or even upper and lower bounds of quantities you don't know.

try to make good guesses, and keep track of these guesses, as they will set bounds on the fidelity of your estimate

- (4). Use ratios when possible by comparing the value of one quantity

 (e.g., force, energy, etc.) in comparison to a related quantity in order to
 eliminate unknown parameters and get a dimensionless parameter
- (5). If possible, exploit plausible scaling behavior of some quantity,
 i.e., estimate an unknown quantity by assuming it scales linearly from known values with some parameter



Making Estimates in Research: How?

More guidelines for making order-of-magnitude estimations:

Checking your estimates:

- (1). Make sure that your estimates and calculations are dimensionally correct! \Rightarrow This is a very powerful tool!
- (2). Check the plausibility of your estimate, if possible e.g., if your answer exceeds the speed of light or the size of the universe, you've got a problem!
- (3). Check the plausibility of your estimate using an alternate calculation method do the two methods agree to within an order of magnitude?
- (4). Perform a "reality check" on your estimate based on the number and size of the approximations you made
- (5). More quantitatively place "bounds" on your estimate:
 To obtain an "upper bound" in equations, put largest estimated values of quantities in the numerator and the smallest estimated values in the denominator

To obtain a "lower bound" – in equations, put smallest estimated values of quantities in the numerator and the largest estimated values in the denominator



Chemistry Estimates: Atoms in a Grain

How many atoms in a grain of sand?



(1). What do we need to know? Size of a grain of sand, size of an atom

Dimension of a grain of sand: assume $L_{sand} \sim 1 \text{ mm} = 10^{-3} \text{ m} \times 10^{10} \text{ Å/m} = 10^7 \text{ Å}$

Dimension of an atom: assume $L_{atom} \sim 1 \text{ Å}$

(2). Do we need a model? Don't worry about geometry...assume a cubic grain of sand!

Volume of grain of sand: $V_{sand} \sim (10^7)^3 \text{ Å}^3 \sim 10^{21} \text{ Å}^3$

Volume of an atom: $V_{atom} \sim (1)^3 \text{ Å}^3 \sim 1 \text{ Å}^3$

(3). Make your estimate!

of atoms in 1 grain ~ (volume of 1 grain)/(volume of 1 atom) ~ 10^{21} atoms per grain

(4). Now, check: How good were our approximations?

Close to N_A... so this is reasonable

"Musical" Estimates: Piano Tuners in CU

How many piano tuners are there in Champaign? (Similar to an original Fermi problem!)

(1). What do we need to know? How many families are there in Champaign? How many families own a piano? How often are pianos tuned? How many tuners are needed?

of people in CU: Estimate 225,000

of families in CU: Estimate 225,000/4 ~ 56,000

of families owning a piano: Estimate 1 in $10 \Rightarrow \sim 6,000$ pianos in Champaign

rate at which pianos are tuned: Estimate 1 time each year $\Rightarrow \sim 6,000$ tunings/year

rate at which piano tuners can perform a tuning: Estimate ~ 4 tunings/day, working ~ 200 days/year (exclude weekends and holidays) \Rightarrow ~ 800 tunings/year per tuner

(2). Make your estimate!

of tuners ~ (rate at which piano tunings are needed)/(rate at which each tuner performs tunings)

 \Rightarrow (6000 tunings/year)/(800 tunings/year*tuner) ~ 8 tuners (± 3 tuners)



"Let me through - I'm a piano tuner!"



From C/U Yellow Pages...

Estimate: ~8 Actual: 7

Culinary Estimates: Supersize Me

How many McDonalds franchises are there in the US?

(1). What do we know? # of McDonalds in Champaign, population of Champaign, population of US

of McDonalds in Champaign: estimate N_{champaign} ~ 7

population of Champaign: ~ 225,000

population of US: ~ 325,000,000

 \Rightarrow Reasonably assume that # of McDonalds franchises scales with population!

(2). Make your estimate!
 Assume simple scaling relationship
 N_{USA} ~ (population of US/population of Champaign)*(# of McDonalds in Champaign)

N_{USA} ~ 10,000 Actual: 13,673



Making Estimates: Hairs on a Human Head

How many hairs on a human head?

(1). What do we need to know? Size of a typical scalp, and approximate number of hairs per square inch

Hairs per inch: 20 - 40 (guess or measure) $\Rightarrow 400 - 1600$ per inch²

(2). Do we need a model? We don't want a precise, specific answer, so assume "average" head that is a hemisphere (keep it simple!)

Radius of "typical" head: ~ 5 inch (guess or measure) \Rightarrow area of scalp ~ $\frac{1}{2}(4\pi r^2)$ ~ 150 inch² (you can take π ~ 3! Excellent!)

(3). Make your estimate!

Note dimensional consistency!

of hairs ~ area of scalp * hairs/unit area ~ 60,000 – 240,000 hairs

OR N ~ 10^5 hairs

Rounding up is OK, even encouraged! \Rightarrow

(4). Now, check: How good were our approximations?



Other resources on making estimates

- A View From the Back of the Envelope http://www.vendian.org/envelope/
- University of Maryland Fermi Problems Site http://www.physics.umd.edu/perg/fermi/fermi.htm
- Old Dominion University Fermi Problems Site http://www.physics.odu.edu/~weinstei/wag.html
- Order of Magnitude Astrophysics http://www.astronomy.ohio-state.edu/~dhw/Oom/questions.html
- Back-of-the-Envelope Physics, Clifford Swartz (Baltimore, Johns Hopkins University Press, 2003).
- The Back of the Envelope, E.M. Purcell, monthly column in the American Journal of Physics, July 1984 – Jan. 1993.
- Consider a Spherical Cow : A Course in Environmental Problem Solving, John Harte (Berkeley, University Science Books, 1988).
- Powers of Ten : About the Relative Size of Things in the Universe, and the Effect of Adding Another Zero, Philip Morrison and Phylis Morrison (Scientific American Library, 1982, 1994).