


Probability and Statistics for Computer Science



Principal Component Analysis ---
Exploring the data in less
dimensions

Credit: wikipedia

Last time

- ✱ Review of Bayesian inference
- ✱ Visualizing high dimensional data & Summarizing data
- ✱ The covariance matrix

Objectives

- ✱ Principal Component Analysis
- ✱ Examples of PCA

Diagonalization of a symmetric matrix

- ✱ If A is an $n \times n$ symmetric square matrix, the eigenvalues are real.
- ✱ If the eigenvalues are also distinct, their eigenvectors are orthogonal
- ✱ We can then scale the eigenvectors to unit length, and place them into an orthogonal matrix $U = [\mathbf{u}_1 \ \mathbf{u}_2 \ \dots \ \mathbf{u}_n]$
- ✱ We can write the diagonal matrix $\Lambda = U^T A U$ such that the diagonal entries of Λ are $\lambda_1, \lambda_2, \dots, \lambda_n$ in that order.

Diagonalization example

✱ For

$$A = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix}$$

Covariance for a pair of components in a data set

- ✱ For the j th and k th components of a data set $\{x\}$

$$\text{cov}(\{x\}; j, k) = \frac{\sum_i (x_i^{(j)} - \text{mean}(\{x^{(j)}\}))(x_i^{(k)} - \text{mean}(\{x^{(k)}\}))^T}{N}$$

Covariance matrix

Data set $\{\mathbf{x}\}$ 7×8

$cov(\{\mathbf{x}\}; 3, 5)$

	1	2	3	4	5	6	7	8
1	*	*	*	*	*	*	*	*
2	*	*	*	*	*	*	*	*
3	*	*	*	*	*	*	*	*
4	*	*	*	*	*	*	*	*
5	*	*	*	*	*	*	*	*
6	*	*	*	*	*	*	*	*
7	*	*	*	*	*	*	*	*

Covmat($\{\mathbf{x}\}$) 7×7

	1	2	3	4	5	6	7
1	*	*	*	*	*	*	*
2	*	*	*	*	*	*	*
3	*	*	*	*	*	*	*
4	*	*	*	*	*	*	*
5	*	*	*	*	*	*	*
6	*	*	*	*	*	*	*
7	*	*	*	*	*	*	*

Properties of Covariance matrix

$$\text{cov}(\{x\}; j, j) = \text{var}(\{x^{(j)}\}) \quad \text{Covmat}(\{\mathbf{x}\}) \quad 7 \times 7$$

- ✱ The diagonal elements of the covariance matrix are just variances of each j th components
- ✱ The off diagonals are covariance between different components

	1	2	3	4	5	6	7
1	*	*	*	*	*	*	*
2	*	*	*	*	*	*	*
3	*	*	*	*	*	*	*
4	*	*	*	*	*	*	*
5	*	*	*	*	*	*	*
6	*	*	*	*	*	*	*
7	*	*	*	*	*	*	*

Properties of Covariance matrix

$$\text{cov}(\{x\}; j, k) = \text{cov}(\{x\}; k, j) \quad \text{Covmat}(\{\mathbf{x}\}) \quad 7 \times 7$$

- ✱ The covariance matrix is **symmetric!**
- ✱ And it's **positive semi-definite**, that is all $\lambda_i \geq 0$
- ✱ Covariance matrix is diagonalizable

	1	2	3	4	5	6	7
1	*	*	*	*	*	*	*
2	*	*	*	*	*	*	*
3	*	*	*	*	*	*	*
4	*	*	*	*	*	*	*
5	*	*	*	*	*	*	*
6	*	*	*	*	*	*	*
7	*	*	*	*	*	*	*

Properties of Covariance matrix

- ✱ If we define X_c as the mean centered matrix for dataset $\{x\}$

$$\text{Covmat}(\{x\}) = \frac{X_c X_c^T}{N}$$

- ✱ The covariance matrix is a $d \times d$ matrix

Covmat($\{\mathbf{x}\}$) 7×7

	1	2	3	4	5	6	7
1	*	*	*	*	*	*	*
2	*	*	*	*	*	*	*
3	*	*	*	*	*	*	*
4	*	*	*	*	*	*	*
5	*	*	*	*	*	*	*
6	*	*	*	*	*	*	*
7	*	*	*	*	*	*	*

$d = 7$

Example: covariance matrix of a data set

(I)

$$A_0 = \begin{bmatrix} 5 & 4 & 3 & 2 & 1 \\ -1 & 1 & 0 & 1 & -1 \end{bmatrix} \begin{matrix} x^{(1)} \\ x^{(2)} \end{matrix}$$

What are the dimensions of the covariance matrix of this data?

- A) 2 by 2
- B) 5 by 5
- C) 5 by 2
- D) 2 by 5

Example: covariance matrix of a data set

(I) Mean centering

$$A_0 = \begin{bmatrix} 5 & 4 & 3 & 2 & 1 \\ -1 & 1 & 0 & 1 & -1 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} 2 & 1 & 0 & -1 & -2 \\ -1 & 1 & 0 & 1 & -1 \end{bmatrix}$$

(II) $A_2 = A_1 A_1^T$

Inner product of each pairs:

$$A_2[1,1] = 10$$

$$A_2[2,2] = 4$$

$$A_2[1,2] = 0$$


(III)

Divide the matrix with N – the number of data points

$$\text{Covmat}(\{\mathbf{x}\}) = \frac{1}{N} A_2 = \frac{1}{5} \begin{bmatrix} 10 & 0 \\ 0 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 0.8 \end{bmatrix}$$

What do the data look like when $\text{Covmat}(\{\mathbf{x}\})$ is diagonal?

$$A_0 = \begin{bmatrix} 5 & 4 & 3 & 2 & 1 \\ -1 & 1 & 0 & 1 & -1 \end{bmatrix} \begin{matrix} x^{(1)} \\ x^{(2)} \end{matrix}$$



$$\text{Covmat}(\{\mathbf{x}\}) = \frac{1}{N} A_2 = \frac{1}{5} \begin{bmatrix} 10 & 0 \\ 0 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 0.8 \end{bmatrix}$$

What is the correlation between the 2 components for the data \mathbf{m} ?

$$\text{Covmat}(\mathbf{m}) = \begin{bmatrix} 20 & 25 \\ 25 & 40 \end{bmatrix}$$

Q. Is this true?

Transforming a matrix with orthonormal matrix only rotates the data


A. Yes

B. No

Dimension Reduction


- ✱ In stead of showing more dimensions through visualization, it's a good idea to do dimension reduction in order to see the major features of the data set.
- ✱ For example, principal component analysis help find the major components of the data set.
- ✱ PCA is essentially about finding eigenvectors of the covariance matrix of the data set $\{x\}$

Dimension reduction from 2D to 1D




Credit: Prof. Forsyth


Step 1: subtract the mean



Step 2: Rotate so that the new data has diagonalized covariance matrix



Step 3: Drop component(s)



Principal Components

- ✱ The columns of U are the normalized eigenvectors of the $\text{Covmat}(\{x\})$ and are called the **principal components** of the data $\{x\}$

Principal components analysis

- * We reduce the dimensionality of dataset $\{\mathbf{x}\}$ represented by matrix $\mathbf{D}_{d \times n}$ from d to s ($s < d$).
- * Step 1. define matrix $\mathbf{m}_{d \times n}$ such that $\mathbf{m} = \mathbf{D} - \text{mean}(\mathbf{D})$
- * Step 2. define matrix $\mathbf{r}_{d \times n}$ such that $\mathbf{r}_i = \mathbf{U}^T \mathbf{m}_i$

Where \mathbf{U}^T satisfies $\mathbf{\Lambda} = \mathbf{U}^T \text{Covmat}(\{\mathbf{x}\}) \mathbf{U}$, $\mathbf{\Lambda}$ is the diagonalization of $\text{Covmat}(\{\mathbf{x}\})$ with the eigenvalues sorted in decreasing order, \mathbf{U} is the orthonormal eigenvectors' matrix

- * Step 3. Define matrix $\mathbf{p}_{d \times n}$ such that \mathbf{p} is \mathbf{r} with the last $d-s$ components of \mathbf{r} made zero.

What happened to the mean?

✱ Step 1.

$$\text{mean}(\mathbf{m}) = \text{mean}(\mathbf{D} - \text{mean}(\mathbf{D})) = 0$$

✱ Step 2.

$$\text{mean}(\mathbf{r}) = \mathbf{U}^T \text{mean}(\mathbf{m}) = \mathbf{U}^T \mathbf{0} = 0$$

✱ Step 3.

$$\text{mean}(\mathbf{p}_i) = \text{mean}(\mathbf{r}_i) = 0 \quad \text{while } i \in 1 : s$$

$$\text{mean}(\mathbf{p}_i) = 0 \quad \text{while } i \in s + 1 : d$$

What happened to the covariances?

✱ Step 1.

$$\text{Covmat}(\mathbf{m}) = \text{Covmat}(\mathbf{D}) = \text{Covmat}(\{\mathbf{x}\})$$

✱ Step 2.

$$\text{Covmat}(\mathbf{r}) = \mathbf{U}^T \text{Covmat}(\mathbf{m}) \mathbf{U} = \mathbf{\Lambda}$$

✱ Step 3. $\text{Covmat}(\mathbf{p})$ is $\mathbf{\Lambda}$ with the last/smallest d -s diagonal terms turned to 0.

Sample covariance matrix

- ✱ In many statistical programs, the sample covariance matrix is defined to be

$$\mathit{Covmat}(\mathbf{m}) = \frac{\mathbf{m} \mathbf{m}^T}{N - 1}$$

- ✱ Similar to what happens to the unbiased standard deviation

PCA an example

✱ Step 1.

$$D = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix} \Rightarrow \text{mean}(\mathbf{D}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{m} = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix}$$

✱ Step 2.

✱ Step 3.

PCA an example

✱ Step 1.

$$D = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix} \Rightarrow \text{mean}(\mathbf{D}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{m} = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix}$$

✱ Step 2.

$$\text{Covmat}(\mathbf{m}) = \begin{bmatrix} 20 & 25 \\ 25 & 40 \end{bmatrix} \Rightarrow \lambda_1 \simeq 57; \quad \lambda_2 \simeq 3$$

$$\Rightarrow \mathbf{U} = \begin{bmatrix} 0.5606288 & -0.8280672 \\ 0.8280672 & 0.5606288 \end{bmatrix} \quad \mathbf{U}^T = \begin{bmatrix} 0.5606288 & 0.8280672 \\ -0.8280672 & 0.5606288 \end{bmatrix}$$

✱ Step 3.

PCA an example

✱ Step 1.

$$D = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix} \Rightarrow \text{mean}(\mathbf{D}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{m} = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix}$$

✱ Step 2.

$$\text{Covmat}(\mathbf{m}) = \begin{bmatrix} 20 & 25 \\ 25 & 40 \end{bmatrix} \Rightarrow \lambda_1 \simeq 57; \quad \lambda_2 \simeq 3$$

$$\Rightarrow \mathbf{U} = \begin{bmatrix} 0.5606288 & -0.8280672 \\ 0.8280672 & 0.5606288 \end{bmatrix} \quad \mathbf{U}^T = \begin{bmatrix} 0.5606288 & 0.8280672 \\ -0.8280672 & 0.5606288 \end{bmatrix}$$

$$\Rightarrow \mathbf{r} = \mathbf{U}^T \mathbf{m} = \begin{bmatrix} 7.478 & -7.211 & 10.549 & -0.267 & -3.071 & -7.478 \\ 1.440 & -0.052 & -1.311 & -1.389 & 2.752 & -1.440 \end{bmatrix}$$

✱ Step 3.

PCA an example

✱ Step 1.

$$D = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix} \Rightarrow \text{mean}(\mathbf{D}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{m} = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix}$$

✱ Step 2.

$$\text{Covmat}(\mathbf{m}) = \begin{bmatrix} 20 & 25 \\ 25 & 40 \end{bmatrix} \Rightarrow \lambda_1 \simeq 57; \quad \lambda_2 \simeq 3$$

$$\Rightarrow \mathbf{U} = \begin{bmatrix} 0.5606288 & -0.8280672 \\ 0.8280672 & 0.5606288 \end{bmatrix} \quad \mathbf{U}^T = \begin{bmatrix} 0.5606288 & 0.8280672 \\ -0.8280672 & 0.5606288 \end{bmatrix}$$

$$\Rightarrow \mathbf{r} = \mathbf{U}^T \mathbf{m} = \begin{bmatrix} 7.478 & -7.211 & 10.549 & -0.267 & -3.071 & -7.478 \\ 1.440 & -0.052 & -1.311 & -1.389 & 2.752 & -1.440 \end{bmatrix}$$

✱ Step 3. $\Rightarrow \mathbf{p} = \begin{bmatrix} 7.478 & -7.211 & 10.549 & -0.267 & -3.071 & -7.478 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

What is this matrix for the previous example?

$$U^T Covmat(\mathbf{m})U = ?$$

What is this matrix for the previous example?

$$U^T Covmat(\mathbf{m})U = ?$$

$$\begin{bmatrix} 57 & 0 \\ 0 & 3 \end{bmatrix}$$

The Mean square error of the projection

- ✱ The mean square error is the sum of the smallest $d-s$ eigenvalues in Λ

$$\frac{1}{N-1} \sum_i \|r_i - p_i\|^2 = \frac{1}{N-1} \sum_i \sum_{j=s+1}^d (r_i^{(j)})^2$$

The Mean square error of the projection

- ✱ The mean square error is the sum of the smallest $d-s$ eigenvalues in Λ

$$\frac{1}{N-1} \sum_i \|r_i - p_i\|^2 = \frac{1}{N-1} \sum_i \sum_{j=s+1}^d (r_i^{(j)})^2 = \sum_{j=s+1}^d \sum_i \frac{1}{N-1} (r_i^{(j)})^2$$

The Mean square error of the projection

- ✱ The mean square error is the sum of the smallest $d-s$ eigenvalues in Λ

$$\begin{aligned} \frac{1}{N-1} \sum_i \|r_i - p_i\|^2 &= \frac{1}{N-1} \sum_i \sum_{j=s+1}^d (r_i^{(j)})^2 = \sum_{j=s+1}^d \sum_i \frac{1}{N-1} (r_i^{(j)})^2 \\ &= \sum_{j=s+1}^d \text{var}(r_i^{(j)}) \end{aligned}$$

The Mean square error of the projection


- ✱ The mean square error is the sum of the smallest $d-s$ eigenvalues in Λ

$$\begin{aligned}\frac{1}{N-1} \sum_i \|r_i - p_i\|^2 &= \frac{1}{N-1} \sum_i \sum_{j=s+1}^d (r_i^{(j)})^2 = \sum_{j=s+1}^d \sum_i \frac{1}{N-1} (r_i^{(j)})^2 \\ &= \sum_{j=s+1}^d \text{var}(r_i^{(j)}) \\ &= \sum_{j=s+1}^d \lambda_j\end{aligned}$$


Examples: Immune Cell Data

- ✱ There are 38816 white blood immune cells from a mouse sample
- ✱ Each immune cell has 40+ features/components
- ✱ Four features are used as illustration.
- ✱ There are at least 3 cell types involved


T cells



B cells




Natural killer cells



Scatter matrix of Immune Cells


- ✱ There are 38816 white blood immune cells from a mouse sample
- ✱ Each immune cell has 40+ features/components
- ✱ Four features are used for the illustration.
- ✱ There are at least 3 cell types involved



PCA of Immune Cells


```
> res1
$values Eigenvalues
[1] 4.7642829 2.1486896 1.3730662
0.4968255

Eigenvectors
$vector
      [,1]  [,2]  [,3]  [,4]
[1,] 0.2476698 0.00801294 -0.6822740
0.6878210
[2,] 0.3389872 -0.72010997 -0.3691532 -
0.4798492
[3,] -0.8298232 0.01550840 -0.5156117 -
0.2128324
[4,] 0.3676152 0.69364033 -0.3638306 -
0.5013477
```



New coordinates in PCA

```
> head(new_coord_t)
      PC1   PC2   PC3   PC4
1  3.6739228  0.1127233 -1.32744266
0.61005994
2 -0.9255199 -2.1016573 -0.80762548 -
0.29104900
3  3.1150230  0.3526459 -0.83994064
0.46074556
4  3.1801414  0.5679807 -0.07097689
0.01539266
5  2.7972723 -0.1073053 -0.39168826 -
0.03981390
6  3.3012610  0.1979659  0.17965423
0.45373049CD3e -0.3676152  0.69364033 -
0.3638306 -0.5013477 [4,]  0.3676152
0.69364033 -0.3638306 -0.5013477
```



What is the percentage of variance that PC₁ covers?

Given the eigenvalues: 4.7642829 2.1486896
1.3730662 0.4968255, what is the
percentage that PC₁ covers?

- A. 54%
- B. 16%
- C. 25%

Reconstructing the data

- ✱ Given the projected data $\mathbf{p}_{d \times n}$ and $\text{mean}(\{\mathbf{x}\})$, we can approximately reconstruct the original data

$$\hat{\mathbf{D}} = \mathbf{U}\mathbf{p} + \text{mean}(\{\mathbf{x}\})$$

- ✱ Each reconstructed data item $\hat{\mathbf{D}}_i$ is a linear combination of the columns of \mathbf{U} weighted by \mathbf{p}_i
- ✱ The columns of \mathbf{U} are the normalized eigenvectors of the $\text{Covmat}(\{\mathbf{x}\})$ and are called the **principal components** of the data $\{\mathbf{x}\}$

End-to-end mean square error

- ✱ Each \mathbf{x}_i becomes \mathbf{r}_i by translation and rotation
- ✱ Each \mathbf{p}_i becomes $\hat{\mathbf{x}}_i$ by the opposite rotation and translation
- ✱ Therefore the end to end mean square error is:
$$\frac{1}{N-1} \sum_i \|\hat{\mathbf{x}}_i - \mathbf{x}_i\|^2 = \frac{1}{N-1} \sum_i \|\mathbf{r}_i - \mathbf{p}_i\|^2 = \sum_{j=s+1}^d \lambda_j$$
- ✱ $\lambda_{s+1}, \dots, \lambda_d$ are the smallest $d-s$ eigenvalues of the $\text{Covmat}(\{\mathbf{x}\})$



PCA: Human face data

- ✱ The dataset consists of 213 images
- ✱ Each image is grayscale and has 64 by 64 resolution
- ✱ We can treat each image as a vector with dimension $d = 4096$



Credit: Prof. Forsyth

How quickly the eigenvalues decrease?




What do the principal components of the images look like?



Mean image

The first 16
principal
components
arranged into
images



Credit: Prof. Forsyth

Reconstruction of the image



The original

1st row show the reconstructions using
some number of principal components
2nd row show the corresponding errors

Mean

1

5

10

20

50

100



Credit: Prof. Forsyth

Q. Which are true?

- A . PCA allows us to project data to the direction along which the data has the biggest variance
- B. PCA allows us to compress data
- C. PCA uses linear transformation to show patterns of data
- D. PCA allows us to visualize data in lower dimensions
- E. All of the above

Assignments

- ✱ Read Chapter 10 of the textbook
- ✱ Week 10 module
- ✱ Next time: Intro to classification

Additional References

- ✱ Robert V. Hogg, Elliot A. Tanis and Dale L. Zimmerman. “Probability and Statistical Inference”
- ✱ Morris H. Degroot and Mark J. Schervish
"Probability and Statistics"

See you next time

*See
You!*

