


# Probability and Statistics for Computer Science



“All models are wrong, but some models are useful” --- George Box

Credit: wikipedia


# Last time

- ✱ Stochastic Gradient Descent
- ✱ Naïve Bayesian Classifier

# Objectives

- ✱ Linear regression
  - ✱ The problem
  - ✱ The least square solution
  - ✱ The training and prediction
  - ✱ The R-squared for the evaluation of the fit.

# Some popular topics in Ngram



# Regression models are Machine learning methods

- ✱ Regression models have been around for a while
- ✱ Dr. Kelvin Murphy's Machine Learning book has 3+ chapters on regression

Wait, have we seen the linear regression before?



# It's about *Relationship* between data features

- ✱ Example: Is the height of people related to their weight?

IDNO	BODYFAT	DENSITY	AGE	WEIGHT	HEIGHT
1	12.6	1.0708	23	154.25	67.75
2	6.9	1.0853	22	173.25	72.25
3	24.6	1.0414	22	154.00	66.25
4	10.9	1.0751	26	184.75	72.25
5	27.8	1.0340	24	184.25	71.25
6	20.6	1.0502	24	210.25	74.75
7	19.0	1.0549	26	181.00	69.75
8	12.8	1.0704	25	176.00	72.50
9	5.1	1.0900	25	191.00	74.00
10	12.0	1.0722	23	198.25	73.50

- ✱  $x$  : HEIGHT,  $y$ : WEIGHT

# Chicago social economic census

- ✱ The census included 77 communities in Chicago
- ✱ The census evaluated the average hardship index of the residents
- ✱ The census evaluated the following parameters for each community:
  - ✱ PERCENT\_OF\_HOUSING\_CROWDED
  - ✱ PERCENT\_HOUSEHOLD\_BELOW\_POVERTY
  - ✱ PERCENT\_AGED\_16p\_UNEMPLOYED
  - ✱ PERCENT\_AGED\_25p\_WITHOUT\_HIGH\_SCHOOL\_DIPLOMA
  - ✱ PERCENT\_AGED\_UNDER\_18\_OR\_OVER\_64
  - ✱ PER\_CAPITA\_INCOME

*Given a new community and its parameters,  
can you predict its average hardship index with all these parameters?*



# The regression problem



# Some terminology

- ✱ Suppose the dataset  $\{(\mathbf{x}, y)\}$  consists of  $N$  labeled items  $(\mathbf{x}_i, y_i)$
- ✱ If we represent the dataset as a table
  - ✱ The  $d$  columns representing  $\{\mathbf{x}\}$  are called **explanatory variables**  $\mathbf{x}^{(j)}$
  - ✱ The numerical column  $y$  is called the **dependent variable**

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

# Variables of the Chicago census

- [1] "PERCENT\_OF\_HOUSING\_CROWDED"
- [2] "PERCENT\_HOUSEHOLDS\_BELOW\_POVERTY"
- [3] "PERCENT\_AGED\_16p\_UNEMPLOYED"
- [4] "PERCENT\_AGED\_25p\_WITHOUT\_HIGH\_SCHOOL\_DIPLOMA"
- [5] "PERCENT\_AGED\_UNDER\_18\_OR\_OVER\_64"
- [6] "PER\_CAPITA\_INCOME"
- [7] "HardshipIndex"

# Which is the dependent variable in the census example?

- A. "PERCENT\_OF\_HOUSING\_CROWDED"
- B. "PERCENT\_AGED\_25p\_WITHOUT\_HIGH\_SCHOOL\_DIPLOMA"
- C. "HardshipIndex"
- D. "PERCENT\_AGED\_UNDER\_18\_OR\_OVER\_64"

# Linear model

- ✱ We begin by modeling  $y$  as a linear function of  $\mathbf{x}^{(j)}$  plus randomness

$$y = \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \dots + \mathbf{x}^{(d)}\beta_d + \xi$$

Where  $\xi$  is a zero-mean random variable that represents model error

- ✱ In vector notation:

$$y = \mathbf{x}^T \boldsymbol{\beta} + \xi$$

Where  $\boldsymbol{\beta}$  is the  $d$ -dimensional vector of coefficients that we train

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

# Each data item gives an equation

✻ The model:  $y = \mathbf{x}^T \boldsymbol{\beta} + \xi = \mathbf{x}^{(1)} \beta_1 + \mathbf{x}^{(2)} \beta_2 + \xi$

Training data

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

# Which together form a matrix equation

✿ The model  $y = \mathbf{x}^T \boldsymbol{\beta} + \xi = \mathbf{x}^{(1)} \beta_1 + \mathbf{x}^{(2)} \beta_2 + \xi$

Training data

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

$$\begin{bmatrix} 0 \\ 2 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 3 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \end{bmatrix}$$

# Which together form a matrix equation

✿ The model  $y = \mathbf{x}^T \boldsymbol{\beta} + \xi = \mathbf{x}^{(1)} \beta_1 + \mathbf{x}^{(2)} \beta_2 + \xi$

Training data

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

$$\begin{bmatrix} 0 \\ 2 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 2 & 3 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \end{bmatrix}$$

$$\mathbf{y} = \mathbf{X} \cdot \boldsymbol{\beta} + \mathbf{e}$$



Q. What's the dimension of matrix X?

A.  $N \times d$

B.  $d \times N$

C.  $N \times N$

D.  $d \times d$

# Training the model is to choose $\beta$

- ✱ Given a training dataset  $\{(\mathbf{x}, y)\}$ , we want to fit a model  $y = \mathbf{x}^T \beta + \xi$

- ✱ Define  $\mathbf{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_N \end{bmatrix}$  and  $X = \begin{bmatrix} \mathbf{x}_1^T \\ \vdots \\ \mathbf{x}_N^T \end{bmatrix}$  and  $\mathbf{e} = \begin{bmatrix} \xi_1 \\ \vdots \\ \xi_N \end{bmatrix}$

- ✱ To train the model, we need to choose  $\beta$  that makes  $\mathbf{e}$  small in the matrix equation  $\mathbf{y} = X \cdot \beta + \mathbf{e}$

# Training using least squares

- ✱ In the least squares method, we aim to **minimize**  $\|\mathbf{e}\|^2$

$$\|\mathbf{e}\|^2 = \|\mathbf{y} - X\boldsymbol{\beta}\|^2 = (\mathbf{y} - X\boldsymbol{\beta})^T (\mathbf{y} - X\boldsymbol{\beta})$$

- ✱ Differentiating with respect to  $\boldsymbol{\beta}$  and setting to zero

$$X^T X \boldsymbol{\beta} - X^T \mathbf{y} = 0$$

- ✱ If  $X^T X$  is invertible, the least squares estimate of the coefficient is:

$$\hat{\boldsymbol{\beta}} = (X^T X)^{-1} X^T \mathbf{y}$$

# Derivation of least square solution



# Least square solution in the project



# Convex set and convex function

- ✱ If a set is convex, any line connecting two points in the set is completely included in the set





Figure 7.4 (a) Illustration of a convex set. (b) Illustration of a nonconvex set.

- ✱ A convex function:  
the area above the curve is convex  
$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$
- ✱ The least square function is **convex**



# What's the dimension of matrix $X^T X$ ?

- A.  $N \times d$
- B.  $d \times N$
- C.  $N \times N$
- D.  $d \times d$

# Is this statement true?

If the matrix  $\mathbf{X}^T\mathbf{X}$  does NOT have zero valued eigenvalues, it is invertible.

- A. TRUE
- B. FALSE



# Training using least squares example

✱ Model:  $y = \mathbf{x}^T \boldsymbol{\beta} + \xi = \mathbf{x}^{(1)} \beta_1 + \mathbf{x}^{(2)} \beta_2 + \xi$

$$\hat{\boldsymbol{\beta}} = (X^T X)^{-1} X^T \mathbf{y} = \begin{bmatrix} 2 \\ -\frac{1}{3} \end{bmatrix}$$

Training data

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	3	0
2	3	2
3	6	5

$$\begin{aligned} \hat{\beta}_1 &= 2 \\ \hat{\beta}_2 &= -\frac{1}{3} \end{aligned}$$

# Prediction

- ✱ If we train the model coefficients  $\hat{\beta}$ , we can predict  $y_0^p$  from  $\mathbf{x}_0$

$$y_0^p = \mathbf{x}_0^T \hat{\beta}$$

- ✱ In the model  $y = \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi$  with  $\hat{\beta} = \begin{bmatrix} 2 \\ -\frac{1}{3} \end{bmatrix}$

- ✱ The prediction for  $\mathbf{x}_0 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  is  $y_0^p$

- ✱ The prediction for  $\mathbf{x}_0 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is  $y_0^p$

# A linear model with constant offset

- ✱ The problem with the model  $y = \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi$  is:

- ✱ Let's add a constant offset  $\beta_0$  to the model

$$y = \beta_0 + \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi$$

# Training and prediction with constant offset

✱ The model  $y = \beta_0 + \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi = \mathbf{x}^T\boldsymbol{\beta} + \xi$

✱ Training data:

<b>1</b>	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$
1	1	3	0
1	2	3	2
1	3	6	5

$$\begin{bmatrix} 1 & x^{(1)} & x^{(2)} \end{bmatrix}$$

$$\hat{\boldsymbol{\beta}} = (X^T X)^{-1} X^T \mathbf{y} = \begin{bmatrix} -3 \\ 2 \\ \frac{1}{3} \end{bmatrix}$$

✱ For  $\mathbf{x}_0 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

$$y_0^p = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} -3 \\ 2 \\ \frac{1}{3} \end{bmatrix} = -3$$

# Variance of the linear regression model

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

- ✱ The random error is uncorrelated to the least square solution of linear combination of explanatory variables.

# Variance of the linear regression model: proof

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

---

**Proof:**

# Variance of the linear regression model: proof

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

**Proof:**  $y = \mathbf{X} \cdot \hat{\boldsymbol{\beta}} + \mathbf{e}$

$$\text{var}[y] = (1/N)(y - \bar{y})^T (y - \bar{y})$$

$$\text{var}[y] = (1/N)([\mathbf{X}\hat{\boldsymbol{\beta}} - \overline{\mathbf{X}\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \bar{\mathbf{e}}])^T ([\mathbf{X}\hat{\boldsymbol{\beta}} - \overline{\mathbf{X}\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \bar{\mathbf{e}}])$$

# Variance of the linear regression model: proof

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

---

**Proof:**

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \bar{\mathbf{e}}])^T ([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \bar{\mathbf{e}}])$$



# Variance of the linear regression model: proof

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

---

## Proof:

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])^T ([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + 2[\mathbf{e} - \overline{\mathbf{e}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

# Variance of the linear regression model: proof

✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

---

**Proof:**

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])^T ([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + 2[\mathbf{e} - \overline{\mathbf{e}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{Because } \overline{\mathbf{e}} = 0 ; \quad \mathbf{e}^T X\hat{\boldsymbol{\beta}} = 0 ; \quad \mathbf{e}^T \mathbf{1} = 0$$

# Variance of the linear regression model: proof

✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

**Proof:**

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])^T ([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + 2[\mathbf{e} - \overline{\mathbf{e}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

Because  $\overline{\mathbf{e}} = 0$  ;  $\mathbf{e}^T X\hat{\boldsymbol{\beta}} = 0$  and  $\mathbf{e}^T \mathbf{1} = 0$  ← Due to Least square minimized

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

# Variance of the linear regression model: proof

✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

**Proof:**

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])^T ([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + 2[\mathbf{e} - \overline{\mathbf{e}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

Because  $\overline{\mathbf{e}} = 0$  ;  $\mathbf{e}^T X\hat{\boldsymbol{\beta}} = 0$  and  $\mathbf{e}^T \mathbf{1} = 0$  ← Due to Least square minimized

$$\text{var}[y] = (1/N)([X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}]^T [X\hat{\boldsymbol{\beta}} - \overline{X\hat{\boldsymbol{\beta}}}] + [\mathbf{e} - \overline{\mathbf{e}}]^T [\mathbf{e} - \overline{\mathbf{e}}])$$

$$\text{var}[y] = \text{var}[X\hat{\boldsymbol{\beta}}] + \text{var}[\mathbf{e}]$$

# Evaluating models using R-squared

- ✱ The least squares estimate satisfies this property

$$\text{var}(\{y_i\}) = \text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\}) + \text{var}(\{\xi_i\})$$

- ✱ This property gives us an evaluation metric called R-squared

$$R^2 = \frac{\text{var}(\{\mathbf{x}_i^T \hat{\boldsymbol{\beta}}\})}{\text{var}(\{y_i\})}$$

- ✱ We have  $0 \leq R^2 \leq 1$  with a larger value meaning a better fit.

Q: What is R-squared if there is only one explanatory variable in the model?




Q: What is R-squared if there is only one explanatory variable in the model?


R-squared would be **the correlation coefficient squared** (textbook pgs 43-44)

# R-squared examples

Chirp frequency vs temperature in crickets



Heart rate vs temperature in humans





# Comparing our example models

$$y = \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi$$

$$y = \beta_0 + \mathbf{x}^{(1)}\beta_1 + \mathbf{x}^{(2)}\beta_2 + \xi$$

$$\hat{\boldsymbol{\beta}} = \begin{bmatrix} 2 \\ -\frac{1}{3} \end{bmatrix}$$

$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$	$\mathbf{x}^T \hat{\boldsymbol{\beta}}$
1	3	0	1
2	3	2	3
3	6	5	4

$$\hat{\boldsymbol{\beta}} = \begin{bmatrix} -3 \\ 2 \\ \frac{1}{3} \end{bmatrix}$$

<b>1</b>	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$y$	$\mathbf{x}^T \hat{\boldsymbol{\beta}}$
1	1	3	0	0
1	2	3	2	2
1	3	6	5	5

# Linear regression model for the Chicago census data

Call:

```
lm(formula = HardshipIndex ~ ., data = dat)
```

Residuals:

Min	1Q	Median	3Q	Max
-15.7157	-1.9230	0.1301	1.9810	8.6719

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )	
(Intercept)	105.1394	37.3622	2.814	0.006346	**
PERCENT_OF_HOUSING_CROWDED	0.7189	0.2753	2.612	0.011014	*
PERCENT_HOUSEHOLDS_BELOW_POVERTY	0.6665	0.0781	8.534	1.90e-12	***
PERCENT_AGED_16p_UNEMPLOYED	0.8023	0.1350	5.941	9.93e-08	***
PERCENT_AGED_25p_WITHOUT_HIGH_SCHOOL_DIPLOMA	0.7751	0.1063	7.293	3.64e-10	***
PERCENT_AGED_UNDER_18_OR_OVER_64	0.4807	0.1202	3.998	0.000156	***
PER_CAPITA_INCOME	-11.8819	3.1888	-3.726	0.000391	***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1


Residual standard error: 3.9 on 70 degrees of freedom

Multiple R-squared: 0.983, Adjusted R-squared: 0.9815

F-statistic: 673.9 on 6 and 70 DF, p-value: < 2.2e-16

# Residual is normally distributed?

The Q-Q plot of the residuals is roughly normal



# Prediction for another community

[1] "PERCENT\_OF\_HOUSING\_CROWDED"  
[2]"PERCENT\_HOUSEHOLDS\_BELOW\_POVERTY"  
[3] "PERCENT\_AGED\_16p\_UNEMPLOYED"  
[4]"PERCENT\_AGED\_25p\_WITHOUT\_HIGH\_SCHOOL\_DIPLOMA"  
[5]  
"PERCENT\_AGED\_UNDER\_18\_OR\_OVER\_64"  
[6]"PER\_CAPITA\_INCOME"

**4.7**

**19.7**

**12.9**

**19.5**

**33.5**


**Log(28202)**

Predicted hardship index: **41.46038**


Note: maximum of hardship index in the training data is 98, minimum is 1

# The clusters of the Chicago communities: clusters and hardship

## Clusters of community




## Hardship index of communities




# The clusters of the Chicago communities: per capital income and hardship

## PER\_CAPITAL\_INCOME

Heatmap of PER\_CAPITA\_INCOME (log scale)




## Hardship index of communities




# The clusters of the Chicago communities: without diploma and hardship

PERCENT\_AGED\_25p\_WITHOUT  
\_HIGH\_SCHOOL\_DIPLOMA



Hardship index of communities



# Assignments

- ✱ Read Chapter 13 of the textbook
- ✱ Next time: More on linear regression



# Additional References

- ✱ Robert V. Hogg, Elliot A. Tanis and Dale L. Zimmerman. “Probability and Statistical Inference”
- ✱ Kelvin Murphy, “Machine learning, A Probabilistic perspective”

See you next time

*See  
You!*

