


Probability and Statistics for Computer Science



Credit: wikipedia

“It’s straightforward to link a number to the outcome of an experiment. The result is a **Random variable.**” --- Prof. Forsythe

Random variable is a function, it is not the same as in **$X = X + 1$**

Last time

✱ Conditional Probability

- ✱ Review

- ✱ Bayes rule

- ✱ Total probability

- ✱ Independence


Random numbers

- * Amount of money on a bet
- * Age at retirement of a population
- * Rate of vehicles passing by the toll
- * Body temperature of a puppy in its pet clinic
- * Level of the intensity of pain in a toothache

Random variable as vectors

Brain imaging
of Human
emotions

- A) Moral
conflict
- B) Multi-task
- C) Rest




A. McDonald et al. NeuroImage doi:
10.1016/j.neuroimage.2016.10.048

Objectives

- ✱ Random Variable
- ✱ Probability distribution
- ✱ Cumulative distribution
- ✱ Joint probability
- ✱ Independence of random variables

Random variables



Random variables


- ✱ The values of a random variable can be either **discrete**, **continuous** or **mixed**.

Discrete Random variables

- ✱ The range of a discrete random variable is a countable set of real numbers.

Random Variable Example

✱ Number of pairs in a hand of 5 cards



- ✱ Let a single outcome be the hand of 5 cards
- ✱ Each outcome maps to values in the set of numbers $\{0, 1, 2\}$

Random Variable Example

- ✱ **Number of pairs in a hand of 6 cards**
- ✱ Let a single outcome be the hand of 6 cards
- ✱ What is the range of values of this random variable?

Q: Random Variable

- ✱ If we roll a 3-sided fair die, and define random variable U , *such that*



A. $\{-1, 0, 1\}$

B. $\{0, 1\}$

Q: Random Variable

✻ If we roll a 3-sided die, and define random variable X so that X takes on -1, 0, and 1 respectively for outcomes showing side 1, 2 and 3. Then what is the range, the set of numbers that X^2 takes on?

A. $\{-1, 0, 1\}$

B. $\{0, 1\}$

Three important facts of Random variables

- ✱ Random variables have **probability functions**
- ✱ Random variables can be **conditioned** on events or other random variables
- ✱ Random variables have **averages**

Random variables have probability functions

✱ Let X be a random variable

✱ The set of outcomes

is an event with probability

$$P(X = x)$$

X is the random variable

is any unique instance that X takes on

Probability Distribution

- ✱ $P(X = x)$ is called the probability distribution for all possible x
- ✱ $P(X = x)$ is also denoted as $P(x)$ or $p(x)$
- ✱ $P(X = x) \geq 0$ for all values that X can take, and is 0 everywhere else
- ✱ The sum of the probability distribution is 1
$$\sum_x P(x) = 1$$



Cumulative distribution

- ✱ $P(X \leq x)$ is called the cumulative distribution function of X
- ✱ $P(X \leq x)$ is also denoted as $f(x)$
- ✱ $P(X \leq x)$ is a non-decreasing function of x

Probability distribution and cumulative distribution

✱ Give the random variable X ,

$$X(\omega) = \begin{cases} 1 & \text{outcome of } \omega \text{ is head} \\ 0 & \text{outcome of } \omega \text{ is tail} \end{cases}$$



Function of random variables: die example

Roll 4-sided fair die twice.

Define these random variables:

X , the values of 1st roll

Y , the values of 2nd roll

Sum $S = X + Y$

Difference $D = X - Y$

| | | | | | |
|-----|---|---|---|---|-----|
| Y | 4 | | | | |
| | 3 | | | | |
| | 2 | | | | |
| | 1 | | | | |
| | | 1 | 2 | 3 | 4 |
| | | | | | X |

Size of Sample Space = ?

Random variable: die example

Roll 4-sided fair die twice.

$$P(X = 1)$$

$$P(Y \leq 2)$$

$$P(S = 7)$$

$$P(D \leq -1)$$

| | | | | | |
|-----|---|---|---|---|-----|
| Y | 4 | | | | |
| | 3 | | | | |
| | 2 | | | | |
| | 1 | | | | |
| | | 1 | 2 | 3 | 4 |
| | | | | | X |

Size of Sample Space
= ?

Random variable: die example

$$S = X + Y$$

| | | | | | | |
|-----|---|---|---|---|---|-----|
| Y | 4 | 5 | 6 | 7 | 8 | |
| | 3 | 4 | 5 | 6 | 7 | |
| | 2 | 3 | 4 | 5 | 6 | |
| | 1 | 2 | 3 | 4 | 5 | |
| | | 1 | 2 | 3 | 4 | X |

$$D = X - Y$$


| | | | | | | |
|-----|---|----|----|----|---|-----|
| Y | 4 | -3 | -2 | -1 | 0 | |
| | 3 | -2 | -1 | 0 | 1 | |
| | 2 | -1 | 0 | 1 | 2 | |
| | 1 | 0 | 1 | 2 | 3 | |
| | | 1 | 2 | 3 | 4 | X |

$$P(S = 7)$$

$$P(D \leq -1)$$


Probability distribution of the sum of two random variables

- ✱ Give the random variable S in the 4-sided die, whose range is $\{2,3,4,5,6,7,8\}$, probability distribution of S .




Probability distribution of the difference of two random variables

- ✱ Give the random variable $D = X - Y$,
what is the probability distribution of D ?



Conditional Probability

✱ The probability of **A** given **B**



$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(B) \neq 0$$

The “Size” analogy

Credit: Prof. Jeremy Orloff &
Jonathan Bloom

Conditional probability distribution of random variables

- ✱ The conditional probability distribution of X given Y is

$$P(x|y) = \frac{P(x, y)}{P(y)} \quad P(y) \neq 0$$

Conditional probability distribution of random variables

- ✱ The conditional probability distribution of X given Y is

$$P(x|y) = \frac{P(x, y)}{P(y)} \quad P(y) \neq 0$$

- ✱ The joint probability distribution of two random variables \mathbf{X} and \mathbf{Y} is

$$P(\{X = x\} \cap \{Y = y\})$$

$$\sum_x P(x|y) = 1$$

Get the marginal from joint distri.

- ✱ We can recover the individual probability distributions from the joint probability distribution

$$P(x) = \sum_y P(x, y)$$

$$P(y) = \sum_x P(x, y)$$

Joint probabilities sum to 1

- ✱ The sum of the joint probability distribution

$$\sum_y \sum_x P(x, y) = 1$$


Joint Probability Example

- ✱ Tossing a coin twice, we define random variable X and Y for each toss.

$$X(\omega) = \begin{cases} 1 & \text{outcome of } \omega \text{ is head} \\ 0 & \text{outcome of } \omega \text{ is tail} \end{cases}$$

$$Y(\omega) = \begin{cases} 1 & \text{outcome of } \omega \text{ is head} \\ 0 & \text{outcome of } \omega \text{ is tail} \end{cases}$$

Joint probability distribution example



Joint Probability Example

Now we define Sum $\mathbf{S} = X + Y$, Difference $\mathbf{D} = X - Y$. \mathbf{S} takes on values $\{0, 1, 2\}$ and \mathbf{D} takes on values $\{-1, 0, 1\}$

$$X(\omega) = \begin{cases} 1 & \text{outcome of } \omega \text{ is head} \\ 0 & \text{outcome of } \omega \text{ is tail} \end{cases}$$


$$Y(\omega) = \begin{cases} 1 & \text{outcome of } \omega \text{ is head} \\ 0 & \text{outcome of } \omega \text{ is tail} \end{cases}$$

Joint Probability Example

| | | 2 nd toss | | P(s, d) |
|----------------------|-------|----------------------|----|---------|
| | | S | D | |
| 1 st toss | Y = 1 | 2 | 0 | |
| | X = 1 | 1 | 1 | |
| | Y = 0 | 1 | -1 | |
| | X = 0 | 0 | 0 | |

Suppose coin is fair, and the tosses are independent

Joint probability distribution example



Independence of random variables

- ✱ Random variable X and Y are independent if


$$P(x, y) = P(x)P(y) \text{ for all } x \text{ and } y$$

- ✱ In the previous coin toss example
 - ✱ Are X and Y independent?
 - ✱ Are S and D independent?

Joint probability distribution example

| | | | | | | |
|-----------|---|---------------|---------------|-------------------|--|---------------|
| $P(x, y)$ | | 0 | 1 | X | | $P(y)$ |
| | 0 | $\frac{1}{4}$ | $\frac{1}{4}$ | \longrightarrow | | $\frac{1}{2}$ |
| | 1 | $\frac{1}{4}$ | $\frac{1}{4}$ | \longrightarrow | | $\frac{1}{2}$ |
| | | \downarrow | \downarrow | | | |
| $P(x)$ | | $\frac{1}{2}$ | $\frac{1}{2}$ | | | |

Joint probability distribution example



Joint probability distribution example

| $P(s, d)$ | | -1 | 0 | 1 | D | $P(s)$ |
|-----------|---|---------------|---------------|---------------|-----|---------------|
| S | 0 | 0 | $\frac{1}{4}$ | 0 | → | $\frac{1}{4}$ |
| | 1 | $\frac{1}{4}$ | 0 | $\frac{1}{4}$ | → | $\frac{1}{2}$ |
| | 2 | 0 | $\frac{1}{4}$ | 0 | → | $\frac{1}{4}$ |
| $P(d)$ | | $\frac{1}{4}$ | $\frac{1}{2}$ | $\frac{1}{4}$ | | |

$$P(S = 1, D = 0) \neq P(S = 1)P(D = 0)$$

Conditional probability distribution example

$$P(s|d) = \frac{P(s, d)}{P(d)}$$

| | | -1 | 0 | 1 | <i>D</i> |
|----------|---|----|---------------|---|----------|
| <i>S</i> | 0 | 0 | $\frac{1}{2}$ | 0 | |
| | 1 | 1 | 0 | 1 | |
| | 2 | 0 | $\frac{1}{2}$ | 0 | |

Bayes rule for random variable

- ✱ Bayes rule for events generalizes to random variables

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

$$= \frac{P(y|x)P(x)}{\sum_x P(y|x)P(x)}$$

Total Probability

Conditional probability distribution example

$$P(s|d) = \frac{P(s, d)}{P(d)}$$

S

0

1

2

| | -1 | 0 | 1 |
|---|----|---------------|---|
| 0 | 0 | $\frac{1}{2}$ | 0 |
| 1 | 1 | 0 | 1 |
| 2 | 0 | $\frac{1}{2}$ | 0 |

D

Assignments

- ✱ Module Week 4
- ✱ Next time: More random variable, Expectations, Variance

Additional References

- ✱ Charles M. Grinstead and J. Laurie Snell
"Introduction to Probability"
- ✱ Morris H. Degroot and Mark J. Schervish
"Probability and Statistics"

See you next time

*See
You!*

