

# CS 373: Theory of Computation

Gul Agha    Mahesh Viswanathan

University of Illinois, Urbana-Champaign

Fall 2010

# Emptiness Problem

Given a CFG  $G$  with start symbol  $S$ , is  $L(G)$  empty?

# Emptiness Problem

Given a CFG  $G$  with start symbol  $S$ , is  $L(G)$  empty?

**Solution:** Check if the start symbol  $S$  is generating.

# Emptiness Problem

Given a CFG  $G$  with start symbol  $S$ , is  $L(G)$  empty?

**Solution:** Check if the start symbol  $S$  is generating. How long does that take?

# Determining generating symbols

## Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

# Determining generating symbols

## Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

- Both for-loops take  $O(n)$  time where  $n = |G|$ .

# Determining generating symbols

## Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

- Both for-loops take  $O(n)$  time where  $n = |G|$ .
- Each iteration of repeat-until loop discovers a new variable. So number of iterations is  $O(n)$ . And total is  $O(n^2)$ .

# Membership Problem

Given a CFG  $G = (V, \Sigma, R, S)$  in **Chomsky Normal Form**, and a string  $w \in \Sigma^*$ , is  $w \in L(G)$ ?

# Membership Problem

Given a CFG  $G = (V, \Sigma, R, S)$  in **Chomsky Normal Form**, and a string  $w \in \Sigma^*$ , is  $w \in L(G)$ ?  
Central question in parsing.

# “Simple” Solution

- Let  $|w| = n$ . Since  $G$  is in Chomsky Normal Form,  $w$  has a parse tree of size  $2n - 1$  iff  $w \in L(G)$

## “Simple” Solution

- Let  $|w| = n$ . Since  $G$  is in Chomsky Normal Form,  $w$  has a parse tree of size  $2n - 1$  iff  $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for  $w$

## “Simple” Solution

- Let  $|w| = n$ . Since  $G$  is in Chomsky Normal Form,  $w$  has a parse tree of size  $2n - 1$  iff  $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for  $w$
- Number of parse trees of size  $2n - 1$  is  $k^{2n-1}$  where  $k$  is the number of variables in  $G$ . So algorithm is exponential in  $n$ !

## “Simple” Solution

- Let  $|w| = n$ . Since  $G$  is in Chomsky Normal Form,  $w$  has a parse tree of size  $2n - 1$  iff  $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for  $w$
- Number of parse trees of size  $2n - 1$  is  $k^{2n-1}$  where  $k$  is the number of variables in  $G$ . So algorithm is exponential in  $n!$
- We will see an algorithm that runs in  $O(n^3)$  time (the constant will depend on  $k$ ).

# First Ideas

## Notation

Suppose  $w = w_1 w_2 \cdots w_n$ , where  $w_i \in \Sigma$ . Let  $w_{i,j}$  denote the substring of  $w$  starting at position  $i$  of length  $j$ . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

# First Ideas

## Notation

Suppose  $w = w_1 w_2 \cdots w_n$ , where  $w_i \in \Sigma$ . Let  $w_{i,j}$  denote the substring of  $w$  starting at position  $i$  of length  $j$ . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

## Main Idea

For every  $A \in V$ , and every  $i \leq n$ ,  $j \leq n + 1 - i$ , we will determine if  $A \xrightarrow{*} w_{i,j}$ .

# First Ideas

## Notation

Suppose  $w = w_1 w_2 \cdots w_n$ , where  $w_i \in \Sigma$ . Let  $w_{i,j}$  denote the substring of  $w$  starting at position  $i$  of length  $j$ . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

## Main Idea

For every  $A \in V$ , and every  $i \leq n$ ,  $j \leq n + 1 - i$ , we will determine if  $A \xrightarrow{*} w_{i,j}$ .

Now,  $w \in L(G)$  iff  $S \xrightarrow{*} w_{1,n} = w$ ; thus, we will solve the membership problem.

# First Ideas

## Notation

Suppose  $w = w_1 w_2 \cdots w_n$ , where  $w_i \in \Sigma$ . Let  $w_{i,j}$  denote the substring of  $w$  starting at position  $i$  of length  $j$ . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

## Main Idea

For every  $A \in V$ , and every  $i \leq n$ ,  $j \leq n + 1 - i$ , we will determine if  $A \xRightarrow{*} w_{i,j}$ .

Now,  $w \in L(G)$  iff  $S \xRightarrow{*} w_{1,n} = w$ ; thus, we will solve the membership problem.

How do we determine if  $A \xRightarrow{*} w_{i,j}$  for every  $A, i, j$ ?

# Base Case

## Substrings of length 1

### Observation

For any  $A, i$ ,  $A \xRightarrow{*} w_{i,1}$  iff  $A \rightarrow w_{i,1}$  is a rule.

# Base Case

## Substrings of length 1

### Observation

For any  $A, i$ ,  $A \xRightarrow{*} w_{i,1}$  iff  $A \rightarrow w_{i,1}$  is a rule.

- Since  $G$  is in Chomsky Normal Form,  $G$  does not have any  $\epsilon$ -rules, nor any unit rules.

# Base Case

## Substrings of length 1

### Observation

For any  $A, i$ ,  $A \xRightarrow{*} w_{i,1}$  iff  $A \rightarrow w_{i,1}$  is a rule.

- Since  $G$  is in Chomsky Normal Form,  $G$  does not have any  $\epsilon$ -rules, nor any unit rules.

Thus, for each  $A$  and  $i$ , one can determine if  $A \xRightarrow{*} w_{i,1}$ .


# Inductive Step

## Longer substrings

Suppose for every variable  $X$  and every  $w_{i,l}$   
( $l < j$ ) we have determined if  $X \xRightarrow{*} w_{i,l}$

# Inductive Step

## Longer substrings




Suppose for every variable  $X$  and every  $w_{i,l}$  ( $l < j$ ) we have determined if  $X \xRightarrow{*} w_{i,l}$

- $A \xRightarrow{*} w_{i,j}$  iff there are variables  $B$  and  $C$  and some  $k < j$  such that  $A \rightarrow BC$  is a rule, and  $B \xRightarrow{*} w_{i,k}$  and  $C \xRightarrow{*} w_{i+k,j-k}$

# Inductive Step

## Longer substrings



Suppose for every variable  $X$  and every  $w_{i,\ell}$  ( $\ell < j$ ) we have determined if  $X \xRightarrow{*} w_{i,\ell}$

- $A \xRightarrow{*} w_{i,j}$  iff there are variables  $B$  and  $C$  and some  $k < j$  such that  $A \rightarrow BC$  is a rule, and  $B \xRightarrow{*} w_{i,k}$  and  $C \xRightarrow{*} w_{i+k,j-k}$
- Since  $k$  and  $j - k$  are both less than  $j$ , we can inductively determine if  $A \xRightarrow{*} w_{i,j}$ .

# Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ .

Initialize:  $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

**for**  $j = 2$  to  $n$  **do**

**for**  $i = 1$  to  $n - j + 1$  **do**

$X_{i,j} = \emptyset$

**for**  $k = 1$  to  $j - 1$  **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

# Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ .

Initialize:  $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

**for**  $j = 2$  to  $n$  **do**

**for**  $i = 1$  to  $n - j + 1$  **do**

$X_{i,j} = \emptyset$

**for**  $k = 1$  to  $j - 1$  **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

**Correctness:** After each iteration of the outermost loop,  $X_{i,j}$  contains exactly the set of variables  $A$  that can derive  $w_{i,j}$ , for each  $i$ .

# Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ .

Initialize:  $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

**for**  $j = 2$  to  $n$  **do**

**for**  $i = 1$  to  $n - j + 1$  **do**

$X_{i,j} = \emptyset$

**for**  $k = 1$  to  $j - 1$  **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

**Correctness:** After each iteration of the outermost loop,  $X_{i,j}$  contains exactly the set of variables  $A$  that can derive  $w_{i,j}$ , for each  $i$ . Time =  $O(n^3)$ .

# Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let  
 $w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$ :

# Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let

$w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ :

| $j/i$ | 1   | 2      | 3      | 4   | 5      |
|-------|-----|--------|--------|-----|--------|
| 5     |     |        |        |     |        |
| 4     |     |        |        |     |        |
| 3     |     |        |        |     |        |
| 2     |     |        |        |     |        |
| 1     | {B} | {A, C} | {A, C} | {B} | {A, C} |
|       | b   | a      | a      | b   | a      |

## Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let

$w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ :

| $j/i$ | 1          | 2          | 3          | 4          | 5          |
|-------|------------|------------|------------|------------|------------|
| 5     |            |            |            |            |            |
| 4     |            |            |            |            |            |
| 3     |            |            |            |            |            |
| 2     | $\{S, A\}$ | $\{B\}$    | $\{S, C\}$ | $\{S, A\}$ |            |
| 1     | $\{B\}$    | $\{A, C\}$ | $\{A, C\}$ | $\{B\}$    | $\{A, C\}$ |
|       | $b$        | $a$        | $a$        | $b$        | $a$        |

## Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let  
 $w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ :

| $j/i$ | 1           | 2          | 3          | 4          | 5          |
|-------|-------------|------------|------------|------------|------------|
| 5     |             |            |            |            |            |
| 4     |             |            |            |            |            |
| 3     | $\emptyset$ | $\{B\}$    | $\{B\}$    |            |            |
| 2     | $\{S, A\}$  | $\{B\}$    | $\{S, C\}$ | $\{S, A\}$ |            |
| 1     | $\{B\}$     | $\{A, C\}$ | $\{A, C\}$ | $\{B\}$    | $\{A, C\}$ |
|       | $b$         | $a$        | $a$        | $b$        | $a$        |

## Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let  
 $w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ :

| $j/i$ | 1           | 2             | 3          | 4          | 5          |
|-------|-------------|---------------|------------|------------|------------|
| 5     |             |               |            |            |            |
| 4     | $\emptyset$ | $\{S, A, C\}$ |            |            |            |
| 3     | $\emptyset$ | $\{B\}$       | $\{B\}$    |            |            |
| 2     | $\{S, A\}$  | $\{B\}$       | $\{S, C\}$ | $\{S, A\}$ |            |
| 1     | $\{B\}$     | $\{A, C\}$    | $\{A, C\}$ | $\{B\}$    | $\{A, C\}$ |
|       | $b$         | $a$           | $a$        | $b$        | $a$        |

## Example

## Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$  Let  
 $w = baaba$ . The sets  $X_{i,j} = \{A \mid A \xrightarrow{*} w_{i,j}\}$ :

| $j/i$ | 1             | 2             | 3          | 4          | 5          |
|-------|---------------|---------------|------------|------------|------------|
| 5     | $\{S, A, C\}$ |               |            |            |            |
| 4     | $\emptyset$   | $\{S, A, C\}$ |            |            |            |
| 3     | $\emptyset$   | $\{B\}$       | $\{B\}$    |            |            |
| 2     | $\{S, A\}$    | $\{B\}$       | $\{S, C\}$ | $\{S, A\}$ |            |
| 1     | $\{B\}$       | $\{A, C\}$    | $\{A, C\}$ | $\{B\}$    | $\{A, C\}$ |
|       | $b$           | $a$           | $a$        | $b$        | $a$        |

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?
- Is  $L(G_1) \cap L(G_2) = \emptyset$ ?

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?
- Is  $L(G_1) \cap L(G_2) = \emptyset$ ?
- Is  $L(G_1) = L(G_2)$ ?

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?
- Is  $L(G_1) \cap L(G_2) = \emptyset$ ?
- Is  $L(G_1) = L(G_2)$ ?
- Is  $G_1$  ambiguous?

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?
- Is  $L(G_1) \cap L(G_2) = \emptyset$ ?
- Is  $L(G_1) = L(G_2)$ ?
- Is  $G_1$  ambiguous?
- Is  $L(G_1)$  inherently ambiguous?

# More Decision Problems

Given a CFGs  $G_1$  and  $G_2$

- Is  $L(G_1) = \Sigma^*$ ?
- Is  $L(G_1) \cap L(G_2) = \emptyset$ ?
- Is  $L(G_1) = L(G_2)$ ?
- Is  $G_1$  ambiguous?
- Is  $L(G_1)$  inherently ambiguous?

All these problems are undecidable.