

Prove that each of the following languages is *not* regular.

1  $\{0^{2^n} \mid n \geq 0\}$ .

**Solution:**

Let  $F = L = \{0^{2^n} \mid n \geq 0\}$ .

Let  $x$  and  $y$  be arbitrary elements of  $F$ .

Then  $x = 0^{2^i}$  and  $y = 0^{2^j}$  for some non-negative integers  $x$  and  $y$ .

Let  $z = 0^{2^i}$ .

Then  $xz = 0^{2^i}0^{2^i} = 0^{2^{i+1}} \in L$ .

And  $yz = 0^{2^j}0^{2^i} = 0^{2^i+2^j} \notin L$ , because  $i \neq j$

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

**Solution:**

For any non-negative integers  $i \neq j$ , the strings  $0^{2^i}$  and  $0^{2^j}$  are distinguished by the suffix  $0^{2^i}$ , because  $0^{2^i}0^{2^i} = 0^{2^{i+1}} \in L$  but  $0^{2^j}0^{2^i} = 0^{2^{i+j}} \notin L$ . Thus  $L$  itself is an infinite fooling set for  $L$ .

2  $\{0^{2^n}1^n \mid n \geq 0\}$

**Solution:**

Let  $F$  be the language  $0^*$ .

Let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = 0^i$  and  $y = 0^j$  for some non-negative integers  $i \neq j$ .

Let  $z = 0^i1^i$ .

Then  $xz = 0^{2i}1^i \in L$ .

And  $yz = 0^{i+j}1^i \notin L$ , because  $i + j \neq 2i$ .

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

**Solution:**

For all non-negative integers  $i \neq j$ , the strings  $0^i$  and  $0^j$  are distinguished by the suffix  $0^i1^i$ , because  $0^{2i}1^i \in L$  but  $0^{i+j}1^i \notin L$ . Thus, the language  $0^*$  is an infinite fooling set for  $L$ .

### Solution:

For all non-negative integers  $i \neq j$ , the strings  $0^{2i}$  and  $0^{2j}$  are distinguished by the suffix  $1^i$ , because  $0^{2i}1^i \in L$  but  $0^{2j}1^i \notin L$ . Thus, the language  $(00)^*$  is an infinite fooling set for  $L$ .

---

3  $\{0^m1^n \mid m \neq 2n\}$

### Solution:

Let  $F$  be the language  $0^*$ .

Let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = 0^i$  and  $y = 0^j$  for some non-negative integers  $i \neq j$ .

Let  $z = 0^i1^i$ .

Then  $xz = 0^{2i}1^i \notin L$ .

And  $yz = 0^{i+j}1^i \in L$ , because  $i + j \neq 2i$ .

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

### Solution:

For all non-negative integers  $i \neq j$ , the strings  $0^{2i}$  and  $0^{2j}$  are distinguished by the suffix  $1^i$ , because  $0^{2i}1^i \notin L$  but  $0^{2j}1^i \in L$ . Thus, the language  $(00)^*$  is an infinite fooling set for  $L$ .

---

4 Strings over  $\{0, 1\}$  where the number of 0s is exactly twice the number of 1s.

### Solution:

Let  $F$  be the language  $0^*$ .

Let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = 0^i$  and  $y = 0^j$  for some non-negative integers  $i \neq j$ .

Let  $z = 0^i1^i$ .

Then  $xz = 0^{2i}1^i \in L$ .

And  $yz = 0^{i+j}1^i \notin L$ , because  $i + j \neq 2i$ .

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

### Solution:

For all non-negative integers  $i \neq j$ , the strings  $0^{2i}$  and  $0^{2j}$  are distinguished by the suffix  $1^i$ , because  $0^{2i}1^i \in L$  but  $0^{2j}1^i \notin L$ . Thus, the language  $(00)^*$  is an infinite fooling set for  $L$ .

### Solution:

If  $L$  were regular, then the language

$$((0+1)^* \setminus L) \cap 0^*1^* = \{0^m1^n \mid m \neq 2n\}$$

would also be regular, because regular languages are closed under complement and intersection. But we just proved that  $\{0^m1^n \mid m \neq 2n\}$  is not regular in problem 3. [Yes, this proof would be worth full credit, either in homework or on an exam.]

- 5 Strings of properly nested parentheses  $()$ , brackets  $[]$ , and braces  $\{\}$ . For example, the string  $([])\{\}$  is in this language, but the string  $([])$  is not, because the left and right delimiters don't match.

### Solution:

Let  $F$  be the language  $(^*$ .

Let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = (^i$  and  $y = (^j$  for some non-negative integers  $i \neq j$ .

Let  $z = )^i$ .

Then  $xz = (^i)^i \in L$ .

And  $yz = (^j)^i \notin L$ , because  $i \neq j$ .

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

### Solution:

For any non-negative integers  $i \neq j$ , the strings  $(^i$  and  $(^j$  are distinguished by the suffix  $)^i$ , because  $(^i)^i \in L$  but  $(^j)^i \notin L$ . Thus, the language  $(^*$  is an infinite fooling set.

- 6 Strings of the form  $w_1\#w_2\#\dots\#w_n$  for some  $n \geq 2$ , where each substring  $w_i$  is a string in  $\{0,1\}^*$ , and some pair of substrings  $w_i$  and  $w_j$  are equal.

### Solution:

Let  $F$  be the language  $0^*$ .

Let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = 0^i$  and  $y = 0^j$  for some non-negative integers  $i \neq j$ .

Let  $z = \#0^i$ .

Then  $xz = 0^i\#0^i \in L$ .

And  $yz = 0^j\#0^i \notin L$ , because  $i \neq j$ .

Thus,  $F$  is a fooling set for  $L$ .

Because  $F$  is infinite,  $L$  cannot be regular.

## Solution:

For any non-negative integers  $i \neq j$ , the strings  $0^i$  and  $0^j$  are distinguished by the suffix  $\#0^i$ , because  $0^i\#0^i \in L$  but  $0^j\#0^i \notin L$ . Thus, the language  $0^*$  is an infinite fooling set.

---

## Extra problems

---

7  $\{0^{n^2} \mid n \geq 0\}$

## Solution:

Let  $x$  and  $y$  be distinct arbitrary strings in  $L$ .

Without loss of generality,  $x = 0^{i^2}$  and  $y = 0^{j^2}$  for some  $i > j \geq 0$ .

Let  $z = 0^{2i+1}$ .

Then  $xz = 0^{i^2+2i+1} = 0^{(i+1)^2} \in L$

On the other hand,  $yz = 0^{i^2+2j+1} \notin L$ , because  $i^2 < i^2 + 2j + 1 < (i+1)^2$ .

Thus,  $z$  distinguishes  $x$  and  $y$ .

We conclude that  $L$  is an infinite fooling set for  $L$ , so  $L$  cannot be regular.

## Solution:

Let  $x$  and  $y$  be distinct arbitrary strings in  $0^*$ .

Without loss of generality,  $x = 0^i$  and  $y = 0^j$  for some  $i > j \geq 0$ .

Let  $z = 0^{i^2+i+1}$ .

Then  $xz = 0^{i^2+2i+1} = 0^{(i+1)^2} \in L$ .

On the other hand,  $yz = 0^{i^2+i+j+1} \notin L$ , because  $i^2 < i^2 + i + j + 1 < (i+1)^2$ .

Thus,  $z$  distinguishes  $x$  and  $y$ .

We conclude that  $0^*$  is an infinite fooling set for  $L$ , so  $L$  cannot be regular.

## Solution:

Let  $x$  and  $y$  be distinct arbitrary strings in  $0000^*$ .

Without loss of generality,  $x = 0^i$  and  $y = 0^j$  for some  $i > j \geq 3$ .

Let  $z = 0^{i^2-i}$ .

Then  $xz = 0^{i^2} \in L$ .

On the other hand,  $yz = 0^{i^2-i+j} \notin L$ , because

$$(i-1)^2 = i^2 - 2i + 1 < i^2 - i < i^2 - i + j < i^2.$$

(The first inequality requires  $i \geq 2$ , and the second  $j \geq 1$ .)

Thus,  $z$  distinguishes  $x$  and  $y$ .

We conclude that  $0000^*$  is an infinite fooling set for  $L$ , so  $L$  cannot be regular.

---

**8**  $\{w \in (0+1)^* \mid w \text{ is the binary representation of a perfect square}\}$

**Solution:**

We design our fooling set around numbers of the form  $(2^k + 1)^2 = 2^{2k} + 2^{k+1} + 1 = 10^{k-2}10^k1 \in L$ , for any integer  $k \geq 2$ . The argument is somewhat simpler if we further restrict  $k$  to be even.

Let  $F = 1(00)^*1$ , and let  $x$  and  $y$  be arbitrary strings in  $F$ .

Then  $x = 10^{2i-2}1$  and  $y = 10^{2j-2}1$ , for some positive integers  $i \neq j$ .

Without loss of generality, assume  $i < j$ . (Otherwise, swap  $x$  and  $y$ .)

Let  $z = 0^{2i}1$ .

Then  $xz = 10^{2i-2}10^{2i}1$  is the binary representation of  $2^{4i} + 2^{2i+1} + 1 = (2^{2i} + 1)^2$ , and therefore  $xz \in L$ .

On the other hand,  $yz = 10^{2j-2}10^{2i}1$  is the binary representation of  $2^{2i+2j} + 2^{2i+1} + 1$ . Simple algebra gives us the inequalities

$$\begin{aligned}(2^{i+j})^2 &= 2^{2i+2j} \\ &< 2^{2i+2j} + 2^{2i+1} + 1 \\ &< 2^{2(i+j)} + 2^{i+j+1} + 1 \\ &= (2^{i+j} + 1)^2.\end{aligned}$$

So  $2^{2i+2j} + 2^{2i+1} + 1$  lies between two consecutive perfect squares, and thus is not a perfect square, which implies that  $yz \notin L$ .

We conclude that  $F$  is a fooling set for  $L$ . Because  $F$  is infinite,  $L$  cannot be regular.