

Prove that each of the following problems is NP-hard.

- 1. Given an undirected graph *G*, does *G* contain a simple path that visits all but 374 vertices?
- 2. Given an undirected graph *G*, does *G* have a spanning tree in which every node has degree at most 374?
- 3. Given an undirected graph *G*, does *G* have a spanning tree with at most 374 leaves?
- 4. Recall that a 5-coloring of a graph *G* is a function that assigns each vertex of *G* a "color" from the set {0, 1, 2, 3, 4}, such that for any edge *uv*, vertices *u* and *v* are assigned different "colors". A 5-coloring is *careful* if the colors assigned to adjacent vertices are not only distinct, but differ by more than 1 (mod 5). Prove that deciding whether a given graph has a careful 5-coloring is NP-hard. [*Hint: Reduce from the standard* 5COLOR *problem.*]



5. Prove that the following problem is NP-hard: Given an undirected graph *G*, find *any* integer k > 374 such that *G* has a proper coloring with *k* colors but *G* does not have a proper coloring with k - 374 colors.

- 6. To think about later: A *bicoloring* of an undirected graph assigns each vertex a set of *two* colors. There are two types of bicoloring: In a *weak* bicoloring, the endpoints of each edge must use *different* sets of colors; however, these two sets may share one color. In a *strong* bicoloring, the endpoints of each edge must use *distinct* sets of colors; that is, they must use four colors altogether. Every strong bicoloring is also a weak bicoloring.
 - (a) Prove that finding the minimum number of colors in a weak bicoloring of a given graph is NP-hard.
 - (b) Prove that finding the minimum number of colors in a strong bicoloring of a given graph is NP-hard.



Left: A weak bicoloring of a 5-clique with four colors. Right A strong bicoloring of a 5-cycle with five colors.