

Describe recursive backtracking algorithms for the following problems. *Don't worry about running times.*

- 1** Given an array $A[1..n]$ of integers, compute the length of a **longest increasing subsequence**.

Solution:

[#1 of ∞] Add a sentinel value $A[0] = -\infty$. Let $LIS(i, j)$ denote the length of the longest increasing subsequence of $A[j..n]$ where every element is larger than $A[i]$. This function obeys the following recurrence:

$$LIS(i, j) = \begin{cases} 0 & \text{if } j > n \\ LIS(i, j+1) & \text{if } j \leq n \text{ and } A[i] \geq A[j] \\ \max\{LIS(i, j+1), 1 + LIS(j, j+1)\} & \text{otherwise} \end{cases}$$

We need to compute $LIS(0, 1)$.

Solution:

[#2 of ∞] Add a sentinel value $A[n+1] = -\infty$. Let $LIS(i, j)$ denote the length of the longest increasing subsequence of $A[1..j]$ where every element is smaller than $A[j]$. This function obeys the following recurrence:

$$LIS(i, j) = \begin{cases} 0 & \text{if } i < 1 \\ LIS(i-1, j) & \text{if } i \geq 1 \text{ and } A[i] \geq A[j] \\ \max\{LIS(i-1, j), 1 + LIS(i-1, i)\} & \text{otherwise} \end{cases}$$

We need to compute $LIS(n, n+1)$.

Solution:

[#3 of ∞] Let $LIS(i)$ denote the length of the longest increasing subsequence of $A[i..n]$ that begins with $A[i]$. This function obeys the following recurrence:

$$LIS(i) = \begin{cases} 1 & \text{if } A[j] \leq A[i] \text{ for all } j > i \\ 1 + \max\{LIS(j) \mid j > i \text{ and } A[j] > A[i]\} & \text{otherwise} \end{cases}$$

(The first case is actually redundant if we define $\max \emptyset = 0$.) We need to compute $\max_i LIS(i)$.

Solution:

[#4 of ∞] Add a sentinel value $A[0] = -\infty$. Let $LIS(i)$ denote the length of the longest increasing subsequence of $A[i..n]$ that begins with $A[i]$. This function obeys the following recurrence:

$$LIS(i) = \begin{cases} 1 & \text{if } A[j] \leq A[i] \text{ for all } j > i \\ 1 + \max\{LIS(j) \mid j > i \text{ and } A[j] > A[i]\} & \text{otherwise} \end{cases}$$

(The first case is actually redundant if we define $\max \emptyset = 0$.) We need to compute $LIS(0) - 1$; the -1 removes the sentinel $-\infty$ from the start of the subsequence.

Solution:

[#5 of ∞] Add sentinel values $A[0] = -\infty$ and $A[n+1] = \infty$. Let $LIS(j)$ denote the length of the longest increasing subsequence of $A[1..j]$ that ends with $A[j]$. This function obeys the following recurrence:

$$LIS(j) = \begin{cases} 1 & \text{if } j = 0 \\ 1 + \max \{ LIS(i) \mid i < j \text{ and } A[i] < A[j] \} & \text{otherwise} \end{cases}$$

We need to compute $LIS(n+1) - 2$; the -2 removes the sentinels $-\infty$ and ∞ from the subsequence.

2 Given an array $A[1..n]$ of integers, compute the length of a **longest decreasing subsequence**.

Solution:

[one of many] Add a sentinel value $A[0] = \infty$. Let $LDS(i, j)$ denote the length of the longest decreasing subsequence of $A[j..n]$ where every element is smaller than $A[i]$. This function obeys the following recurrence:

$$LDS(i, j) = \begin{cases} 0 & \text{if } j > n \\ LDS(i, j+1) & \text{if } j \leq n \text{ and } A[i] \leq A[j] \\ \max \{ LDS(i, j+1), 1 + LDS(j, j+1) \} & \text{otherwise} \end{cases}$$

We need to compute $LDS(0, 1)$.

Solution:

[clever] Multiply every element of A by -1 , and then compute the length of the longest increasing subsequence using the algorithm from problem 1.

3 Given an array $A[1..n]$ of integers, compute the length of a **longest alternating subsequence**.

Solution:

[one of many] We define two functions:

- Let $LAS^+(i, j)$ denote the length of the longest alternating subsequence of $A[j..n]$ whose first element (if any) is larger than $A[i]$ and whose second element (if any) is smaller than its first.
- Let $LAS^-(i, j)$ denote the length of the longest alternating subsequence of $A[j..n]$ whose first element (if any) is smaller than $A[i]$ and whose second element (if any) is larger than its first.

These two functions satisfy the following mutual recurrences:

$$LAS^+(i, j) = \begin{cases} 0 & \text{if } j > n \\ LAS^+(i, j+1) & \text{if } j \leq n \text{ and } A[j] \leq A[i] \\ \max \{ LAS^+(i, j+1), 1 + LAS^-(j, j+1) \} & \text{otherwise} \end{cases}$$
$$LAS^-(i, j) = \begin{cases} 0 & \text{if } j > n \\ LAS^-(i, j+1) & \text{if } j \leq n \text{ and } A[j] \geq A[i] \\ \max \{ LAS^-(i, j+1), 1 + LAS^+(j, j+1) \} & \text{otherwise} \end{cases}$$

To simplify computation, we consider two different sentinel values $A[0]$. First we set $A[0] = -\infty$ and let $\ell^+ = LAS^+(0, 1)$. Then we set $A[0] = +\infty$ and let $\ell^- = LAS^-(0, 1)$. Finally, the length of the longest alternating subsequence of A is $\max \{ \ell^+, \ell^- \}$.

Solution:

[one of many] We define two functions:

- Let $LAS^+(i)$ denote the length of the longest alternating subsequence of $A[i..n]$ that starts with $A[i]$ and whose second element (if any) is larger than $A[i]$.
- Let $LAS^-(i)$ denote the length of the longest alternating subsequence of $A[i..n]$ that starts with $A[i]$ and whose second element (if any) is smaller than $A[i]$.

These two functions satisfy the following mutual recurrences:

$$LAS^+(i) = \begin{cases} 1 & \text{if } A[j] \leq A[i] \text{ for all } j > i \\ 1 + \max \{ LAS^-(j) \mid j > i \text{ and } A[j] > A[i] \} & \text{otherwise} \end{cases}$$
$$LAS^-(i) = \begin{cases} 1 & \text{if } A[j] \geq A[i] \text{ for all } j > i \\ 1 + \max \{ LAS^+(j) \mid j > i \text{ and } A[j] < A[i] \} & \text{otherwise} \end{cases}$$

We need to compute $\max_i \max \{ LAS^+(i), LAS^-(i) \}$.

To think about later:

- 1 Given an array $A[1..n]$ of integers, compute the length of a longest **convex** subsequence of A .

Solution:

Let $LCS(i, j)$ denote the length of the longest convex subsequence of $A[i..n]$ whose first two elements are $A[i]$ and $A[j]$. This function obeys the following recurrence:

$$LCS(i, j) = 1 + \max \{ LCS(j, k) \mid j < k \leq n \text{ and } A[i] + A[k] > 2A[j] \}$$

Here we define $\max \emptyset = 0$; this gives us a working base case. The length of the longest convex subsequence is $\max_{1 \leq i < j \leq n} LCS(i, j)$.

Solution:

[with sentinels] Assume without loss of generality that $A[i] \geq 0$ for all i . (Otherwise, we can add $|m|$ to each $A[i]$, where m is the smallest element of $A[1..n]$.) Add two sentinel values $A[0] = 2M + 1$ and $A[-1] = 4M + 3$, where M is the largest element of $A[1..n]$.

Let $LCS(i, j)$ denote the length of the longest convex subsequence of $A[i..n]$ whose first two elements are $A[i]$ and $A[j]$. This function obeys the following recurrence:

$$LCS(i, j) = 1 + \max \{ LCS(j, k) \mid j < k \leq n \text{ and } A[i] + A[k] > 2A[j] \}$$

Here we define $\max \emptyset = 0$; this gives us a working base case.

Finally, we claim that the length of the longest convex subsequence of $A[1..n]$ is $LCS(-1, 0) - 2$.

Proof: First, consider any convex subsequence S of $A[1..n]$, and suppose its first element is $A[i]$. Then we have $A[-1] - 2A[0] + A[i] = 4M + 3 - 2(2M + 1) + A[i] = A[i] + 1 > 0$, which implies that $A[-1] \cdot A[0] \cdot S$ is a convex subsequence of $A[-1..n]$. So the longest convex subsequence of $A[1..n]$ has length at most $LCS(-1, 0) - 2$.

On the other hand, removing $A[-1]$ and $A[0]$ from any convex subsequence of $A[-1..n]$ leaves a convex subsequence of $A[1..n]$. So the longest subsequence of $A[1..n]$ has length at least $LCS(-1, 0) - 2$. ■

2 Given an array $A[1..n]$, compute the length of a longest *palindrome* subsequence of A .

Solution:

[naive] Let $LPS(i, j)$ denote the length of the longest palindrome subsequence of $A[i..j]$. This function obeys the following recurrence:

$$LPS(i, j) = \begin{cases} 0 & \text{if } i > j \\ 1 & \text{if } i = j \\ \max \left\{ \begin{array}{l} LPS(i+1, j) \\ LPS(i, j-1) \end{array} \right\} & \text{if } i < j \text{ and } A[i] \neq A[j] \\ \max \left\{ \begin{array}{l} 2 + LPS(i+1, j-1) \\ LPS(i+1, j) \\ LPS(i, j-1) \end{array} \right\} & \text{otherwise} \end{cases}$$

We need to compute $LPS(1, n)$.

Solution:

[with greedy optimization] Let $LPS(i, j)$ denote the length of the longest palindrome subsequence of $A[i..j]$. Before stating a recurrence for this function, we make the following useful observation.

Claim 0.1. *If $i < j$ and $A[i] = A[j]$, then $LPS(i, j) = 2 + LPS(i+1, j-1)$.*

Proof: Suppose $i < j$ and $A[i] = A[j]$. Fix an arbitrary longest palindrome subsequence S of $A[i..j]$. There are four cases to consider.

- If S uses neither $A[i]$ nor $A[j]$, then $A[i] \bullet S \bullet A[j]$ is a palindrome subsequence of $A[i..j]$ that is longer than S , which is impossible.
- Suppose S uses $A[i]$ but not $A[j]$. Let $A[k]$ be the last element of S . If $k = i$, then $A[i] \bullet A[j]$ is a palindrome subsequence of $A[i..j]$ that is longer than S , which is impossible. Otherwise, replacing $A[k]$ with $A[j]$ gives us a palindrome subsequence of $A[i..j]$ with the same length as S that uses both $A[i]$ and $A[j]$.
- Suppose S uses $A[j]$ but not $A[i]$. Let $A[h]$ be the first element of S . If $h = j$, then $A[i] \bullet A[j]$ is a palindrome subsequence of $A[i..j]$ that is longer than S , which is impossible. Otherwise, replacing $A[h]$ with $A[i]$ gives us a palindrome subsequence of $A[i..j]$ with the same length as S that uses both $A[i]$ and $A[j]$.
- Finally, S might include both $A[i]$ and $A[j]$.

In all cases, we find either a contradiction or a longest palindrome subsequence of $A[i..j]$ that uses both $A[i]$ and $A[j]$. ■

Claim 1 implies that the function LPS satisfies the following recurrence:

$$LPS(i, j) = \begin{cases} 0 & \text{if } i > j \\ 1 & \text{if } i = j \\ \max \{ LPS(i+1, j), LPS(i, j-1) \} & \text{if } i < j \text{ and } A[i] \neq A[j] \\ 2 + LPS(i+1, j-1) & \text{otherwise} \end{cases}$$

We need to compute $LPS(1, n)$.