

Prove that each of the following languages is *not* regular.

1. $\{\mathbf{0}^{2n}\mathbf{1}^n \mid n \geq 0\}$

Solution (verbose): Let F be the language $\mathbf{0}^*$.

Let x and y be arbitrary strings in F .

Then $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some non-negative integers $i \neq j$.

Let $z = \mathbf{0}^i\mathbf{1}^i$.

Then $xz = \mathbf{0}^{2i}\mathbf{1}^i \in L$.

And $yz = \mathbf{0}^{i+j}\mathbf{1}^i \notin L$, because $i + j \neq 2i$.

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise): For all non-negative integers $i \neq j$, the strings $\mathbf{0}^i$ and $\mathbf{0}^j$ are distinguished by the suffix $\mathbf{0}^i\mathbf{1}^i$, because $\mathbf{0}^{2i}\mathbf{1}^i \in L$ but $\mathbf{0}^{i+j}\mathbf{1}^i \notin L$. Thus, the language $\mathbf{0}^*$ is an infinite fooling set for L . ■

Solution (concise, different fooling set): For all non-negative integers $i \neq j$, the strings $\mathbf{0}^{2i}$ and $\mathbf{0}^{2j}$ are distinguished by the suffix $\mathbf{1}^i$, because $\mathbf{0}^{2i}\mathbf{1}^i \in L$ but $\mathbf{0}^{2j}\mathbf{1}^i \notin L$. Thus, the language $(\mathbf{00})^*$ is an infinite fooling set for L . ■

2. $\{\mathbf{0}^m \mathbf{1}^n \mid m \neq 2n\}$

Solution (verbose): Let F be the language $\mathbf{0}^*$.

Let x and y be arbitrary strings in F .

Then $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some non-negative integers $i \neq j$.

Let $z = \mathbf{0}^i \mathbf{1}^i$.

Then $xz = \mathbf{0}^{2i} \mathbf{1}^i \notin L$.

And $yz = \mathbf{0}^{i+j} \mathbf{1}^i \in L$, because $i + j \neq 2i$.

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise, different fooling set): For all non-negative integers $i \neq j$, the strings $\mathbf{0}^{2i}$ and $\mathbf{0}^{2j}$ are distinguished by the suffix $\mathbf{1}^i$, because $\mathbf{0}^{2i} \mathbf{1}^i \notin L$ but $\mathbf{0}^{2j} \mathbf{1}^i \in L$. Thus, the language $(\mathbf{00})^*$ is an infinite fooling set for L . ■

3. $\{\mathbf{0}^{2^n} \mid n \geq 0\}$

Solution (verbose): Let $F = L = \{\mathbf{0}^{2^n} \mid n \geq 0\}$.

Let x and y be arbitrary elements of F .

Then $x = \mathbf{0}^{2^i}$ and $y = \mathbf{0}^{2^j}$ for some non-negative integers x and y .

Let $z = \mathbf{0}^{2^i}$.

Then $xz = \mathbf{0}^{2^i} \mathbf{0}^{2^i} = \mathbf{0}^{2^{i+1}} \in L$.

And $yz = \mathbf{0}^{2^j} \mathbf{0}^{2^i} = \mathbf{0}^{2^i+2^j} \notin L$, because $i \neq j$

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise): For any non-negative integers $i \neq j$, the strings $\mathbf{0}^{2^i}$ and $\mathbf{0}^{2^j}$ are distinguished by the suffix $\mathbf{0}^{2^i}$, because $\mathbf{0}^{2^i} \mathbf{0}^{2^i} = \mathbf{0}^{2^{i+1}} \in L$ but $\mathbf{0}^{2^j} \mathbf{0}^{2^i} = \mathbf{0}^{2^i+2^j} \notin L$. Thus L itself is an infinite fooling set for L . ■

4. Strings over $\{\mathbf{0}, \mathbf{1}\}$ where the number of $\mathbf{0}$ s is exactly twice the number of $\mathbf{1}$ s.

Solution (verbose): Let F be the language $\mathbf{0}^*$.

Let x and y be arbitrary strings in F .

Then $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some non-negative integers $i \neq j$.

Let $z = \mathbf{0}^i \mathbf{1}^i$.

Then $xz = \mathbf{0}^{2i} \mathbf{1}^i \in L$.

And $yz = \mathbf{0}^{i+j} \mathbf{1}^i \notin L$, because $i + j \neq 2i$.

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise, different fooling set): For all non-negative integers $i \neq j$, the strings 0^{2i} and 0^{2j} are distinguished by the suffix 1^i , because $0^{2i}1^i \in L$ but $0^{2j}1^i \notin L$. Thus, the language $(00)^*$ is an infinite fooling set for L . ■

Solution (closure properties): If L were regular, then the language

$$L \cap 0^*1^* = \{0^{2n}1^n \mid n \geq 0\}$$

would also be regular since regular languages are closed under intersection but we have seen in Problem 1 that $\{0^{2n}1^n \mid n \geq 0\}$ is not regular.

Another solution based on closure properties. If L were regular, then the language

$$((0+1)^* \setminus L) \cap 0^*1^* = \{0^m1^n \mid m \neq 2n\}$$

would also be regular, because regular languages are closed under complement and intersection. But we just proved that $\{0^m1^n \mid m \neq 2n\}$ is not regular in problem 2. [Yes, this proof would be worth full credit, either in homework or on an exam.]

Note that the proofs based on closure properties relied on non-regularity of some previously known languages. One could also think of the proofs as allowing you to simplify the initial language to a more structured one which may be easier to work with. ■

5. Strings of properly nested parentheses $()$, brackets $[\]$, and braces $\{\}$. For example, the string $([\])\{\}$ is in this language, but the string $([\])$ is not, because the left and right delimiters don't match.

Solution (verbose): Let F be the language $(^*$.

Let x and y be arbitrary strings in F .

Then $x = (^i$ and $y = (^j$ for some non-negative integers $i \neq j$.

Let $z =)^i$.

Then $xz = (^i)^i \in L$.

And $yz = (^j)^i \notin L$, because $i \neq j$.

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise): For any non-negative integers $i \neq j$, the strings $(^i$ and $(^j$ are distinguished by the suffix $)^i$, because $(^i)^i \in L$ but $(^j)^i \notin L$. Thus, the language $(^*$ is an infinite fooling set. ■

Solution (closure properties): If L were regular, then the language $L \cap (^*)^* = \{ (^n)^n \mid n \geq 0 \}$ would be regular. The language $\{ (^n)^n \mid n \geq 0 \}$ is the same as $\{ \mathbf{0}^n \mathbf{1}^n \mid n \geq 0 \}$ modulo changing the symbol names and is not regular from lecture. Thus L is not regular. ■

6. Strings of the form $w_1 \# w_2 \# \dots \# w_n$ for some $n \geq 2$, where each substring w_i is a string in $\{ \mathbf{0}, \mathbf{1} \}^*$, and some pair of substrings w_i and w_j are equal.

Solution (verbose): Let F be the language $\mathbf{0}^*$.

Let x and y be arbitrary strings in F .

Then $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some non-negative integers $i \neq j$.

Let $z = \# \mathbf{0}^i$.

Then $xz = \mathbf{0}^i \# \mathbf{0}^i \in L$.

And $yz = \mathbf{0}^j \# \mathbf{0}^i \notin L$, because $i \neq j$.

Thus, F is a fooling set for L .

Because F is infinite, L cannot be regular. ■

Solution (concise): For any non-negative integers $i \neq j$, the strings $\mathbf{0}^i$ and $\mathbf{0}^j$ are distinguished by the suffix $\# \mathbf{0}^i$, because $\mathbf{0}^i \# \mathbf{0}^i \in L$ but $\mathbf{0}^j \# \mathbf{0}^i \notin L$. Thus, the language $\mathbf{0}^*$ is an infinite fooling set. ■

Work on these later:

7. $\{\mathbf{0}^{n^2} \mid n \geq 0\}$

Solution: Let x and y be distinct arbitrary strings in L .

Without loss of generality, $x = \mathbf{0}^{i^2}$ and $y = \mathbf{0}^{j^2}$ for some $i > j \geq 0$.

Let $z = \mathbf{0}^{2j+1}$.

Then $yz = \mathbf{0}^{j^2+2j+1} = \mathbf{0}^{(j+1)^2} \in L$

On the other hand, $xz = \mathbf{0}^{i^2+2j+1} \notin L$, because $i^2 < i^2 + 2j + 1 < (i + 1)^2$.

Thus, z distinguishes x and y .

We conclude that L is an infinite fooling set for L , so L cannot be regular. ■

Solution: Let x and y be distinct arbitrary strings in $\mathbf{0}^*$.

Without loss of generality, $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some $i > j \geq 0$.

Let $z = \mathbf{0}^{i^2+i+1}$.

Then $xz = \mathbf{0}^{i^2+2i+1} = \mathbf{0}^{(i+1)^2} \in L$.

On the other hand, $yz = \mathbf{0}^{i^2+i+j+1} \notin L$, because $i^2 < i^2 + i + j + 1 < (i + 1)^2$.

Thus, z distinguishes x and y .

We conclude that $\mathbf{0}^*$ is an infinite fooling set for L , so L cannot be regular. ■

Solution: Let x and y be distinct arbitrary strings in $\mathbf{0}\mathbf{0}\mathbf{0}\mathbf{0}^*$.

Without loss of generality, $x = \mathbf{0}^i$ and $y = \mathbf{0}^j$ for some $i > j \geq 3$.

Let $z = \mathbf{0}^{i^2-i}$.

Then $xz = \mathbf{0}^{i^2} \in L$.

On the other hand, $yz = \mathbf{0}^{i^2-i+j} \notin L$, because

$$(i-1)^2 = i^2 - 2i + 1 < i^2 - i < i^2 - i + j < i^2.$$

(The first inequality requires $i \geq 2$, and the second $j \geq 1$.)

Thus, z distinguishes x and y .

We conclude that $\mathbf{0}\mathbf{0}\mathbf{0}\mathbf{0}^*$ is an infinite fooling set for L , so L cannot be regular. ■

8. $\{w \in (\mathbf{0} + \mathbf{1})^* \mid w \text{ is the binary representation of a perfect square}\}$

Solution: We design our fooling set around numbers of the form $(2^k + 1)^2 = 2^{2k} + 2^{k+1} + 1 = \mathbf{10}^{k-2}\mathbf{10}^k\mathbf{1} \in L$, for any integer $k \geq 2$. The argument is somewhat simpler if we further restrict k to be even.

Let $F = \mathbf{1}(\mathbf{00})^*\mathbf{1}$, and let x and y be arbitrary strings in F .

Then $x = \mathbf{10}^{2i-2}\mathbf{1}$ and $y = \mathbf{10}^{2j-2}\mathbf{1}$, for some positive integers $i \neq j$.

Without loss of generality, assume $i < j$. (Otherwise, swap x and y .)

Let $z = \mathbf{0}^{2i}\mathbf{1}$.

Then $xz = \mathbf{10}^{2i-2}\mathbf{10}^{2i}\mathbf{1}$ is the binary representation of $2^{4i} + 2^{2i+1} + 1 = (2^{2i} + 1)^2$, and therefore $xz \in L$.

On the other hand, $yz = \mathbf{10}^{2j-2}\mathbf{10}^{2i}\mathbf{1}$ is the binary representation of $2^{2i+2j} + 2^{2i+1} + 1$. Simple algebra gives us the inequalities

$$\begin{aligned} (2^{i+j})^2 &= 2^{2i+2j} \\ &< 2^{2i+2j} + 2^{2i+1} + 1 \\ &< 2^{2(i+j)} + 2^{i+j+1} + 1 \\ &= (2^{i+j} + 1)^2. \end{aligned}$$

So $2^{2i+2j} + 2^{2i+1} + 1$ lies between two consecutive perfect squares, and thus is not a perfect square, which implies that $yz \notin L$.

We conclude that F is a fooling set for L . Because F is infinite, L cannot be regular. ■