Light and Color



"Empire of Light", Magritte


Computational Photography
Derek Hoiem, University of Illinois

Announcements

- Project 1 due next Mon (9/14), 11:59pm
 - Choosing cutoff: sigma > 1
 - Remember to convert images to double or single
 - Don't use built-in code for pyramids, contrast equalization etc., ask if not sure


Today's class

 How is incoming light measured by the eye or camera?




Today's class

- How is incoming light measured by the eye or camera?
- How is light reflected from a surface?










Slide: Forsyth


The Eye



The human eye is a camera!

- Iris colored annulus with radial muscles
- Pupil the hole (aperture) whose size is controlled by the iris
- What's the "film"?
 - photoreceptor cells (rods and cones) in the retina

Retina up-close




Two types of light-sensitive receptors

Cones


cone-shaped less sensitive operate in high light color vision

Rods


rod-shaped highly sensitive operate at night gray-scale vision slower to respond



Rod / Cone sensitivity




Distribution of Rods and Cones




Night Sky: why are there more stars off-center?

Electromagnetic Spectrum




Human Luminance Sensitivity Function

Visible Light



The Physics of Light


Any patch of light can be completely described physically by its spectrum: the number of photons (per time unit) at each wavelength 400 - 700 nm.




The Physics of Light

Some examples of the spectra of light sources


A. Ruby Laser




B. Gallium Phosphide Crystal



C. Tungsten Lightbulb




D. Normal Daylight




The Physics of Light

Some examples of the <u>reflectance</u> spectra of <u>surfaces</u>




© Stephen E. Palmer, 2002


More Spectra



Physiology of Color Vision

Three kinds of cones:






WAVELENGTH (nm.)


- Why are M and L cones so close?
- Why are there 3?

3 is better than 2...




- "M" and "L" on the X-chromosome
 - Why men are more likely to be color blind (see what it's like: http://www.vischeck.com/vischeck/vischeckURL.php)
- "L" has high variation, so some women are tetrachromatic
- Some animals have 1 (night animals), 2 (e.g., dogs), 4 (fish, birds), 5 (pigeons, some reptiles/amphibians), or even 12 (mantis shrimp)

We don't perceive a spectrum (or even RGB)



- We perceive
 - Hue: mean wavelength, color
 - Saturation: variance, vividness
 - Intensity: total amount of light
- Same perceived color can be recreated with combinations of three primary colors ("trichromacy")

Trichromacy and CIE-XYZ




Perceptual equivalents with RGB

Perceptual equivalents with CIE-XYZ


$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \frac{1}{0.17697} \begin{bmatrix} 0.49 & 0.31 & 0.20 \\ 0.17697 & 0.81240 & 0.01063 \\ 0.00 & 0.01 & 0.99 \end{bmatrix} \begin{bmatrix} R \\ G \\ B \end{bmatrix}$$

CIE-XYZ




$$x = \frac{X}{X + Y + Z}$$

$$y = \frac{Y}{X + Y + Z}$$




RGB portion is in triangle


Color Sensing: Bayer Grid



Estimate RGB at each cell from neighboring values




Alternative to Bayer: RGB+W



How is light reflected from a surface?


Depends on

- Illumination properties: wavelength, orientation, intensity
- Surface properties: material, surface orientation, roughness, etc.



Lambertian surface

- Some light is absorbed (function of albedo)
- Remaining light is reflected in all directions (diffuse reflection)
- Examples: soft cloth, concrete, matte paints



Diffuse reflection

Intensity *does* depend on illumination angle because less light comes in at oblique angles.


 ρ = albedo

S =directional source

N =surface normal

I = image intensity

$$I(x) = \rho(x)(S \cdot N(x))$$




Slide: Forsyth



Diffuse reflection

Perceived intensity does *not* depend on viewer angle.


- Amount of reflected light proportional to cos(theta)
- Visible solid angle also proportional to cos(theta)



http://en.wikipedia.org/wiki/Lambert%27s_cosine_law

Specular Reflection

- Reflected direction depends on light orientation and surface normal
- E.g., mirrors are fully specular







Flickr, by piratejohnny

Many surfaces have both specular and diffuse components

 Specularity = spot where specular reflection dominates (typically reflects light source)







Photo: northcountryhardwoodfloors.com


BRDF: Bidirectional Reflectance Distribution Function

 Model of local reflection that tells how bright a surface appears when viewed from one direction when light falls on it from another




More complicated effects



















Inter-reflection is a major source of light




Inter-reflection affects the apparent color of objects




From Koenderink slides on image texture and the flow of light

The color of objects


- Colored light arriving at the camera involves two effects
 - The color of the light source (illumination + inter-reflections)
 - The color of the surface








Color constancy


- Interpret surface in terms of albedo or "true color", rather than observed intensity
 - Humans are good at it
 - Computers are not nearly as good


















Shadows cast by a point source

- A point that can't see the source is in shadow
- For point sources, the geometry is simple




Slide: Forsyth

Area sources


 Examples: diffuser boxes, white walls

 The energy received at a point due to an area source is obtained by adding up the contribution of small elements over the whole source




Slide: Forsyth

Area Source Shadows



Shading and shadows are major cues to shape and position



From Koenderink slides on image texture and the flow of light

Slide: Forsyth


Recap





- 1. Why is (2) brighter than (1)? Each points to the asphalt.
- 2. Why is (4) darker than (3)? 4 points to the marking.
- 3. Why is (5) brighter than (3)? Each points to the side of the wooden block.
- 4. Why isn't (6) black, given that there is no direct path from it to the sun?


Things to remember

- Light has a spectrum of wavelengths
 - Humans (and RGB cameras) have color sensors sensitive to three ranges
- Observed light depends on: illumination intensities, surface orientation, material (albedo, specular component, diffuse component), etc.
- Every object is an indirect light source for every other
- Shading and shadows are informative about shape and position










Take-home questions



A. For each of the arrows in the above image, name the reasons the pixel near the end of the arrow has its brightness value and explain very briefly. The arrow pointing to milk is pointing to the thin bright line at the edge of the piece of apple; the arrow pointing to the spoon handle is pointing to the bright area on the handle.

Possible factors: albedo, shadows, texture, specularities, curvature, lighting direction