

# CS 473: Fundamental Algorithms, Spring 2013

## Discussion 3

January 29, 2013

### 3.1. 2SAT.

You are given a boolean formula that is a 2CNF. That is, every clause is the OR of two boolean variables, and the formula is the conjunction of the clauses. For an example, consider the following formula:

$$F = (x_1 \vee \bar{x}_2) \wedge (\bar{x}_3 \vee x_4) \wedge (\bar{x}_1 \vee \bar{x}_4) \wedge (\bar{x}_2 \vee \bar{x}_3).$$

- (A) What is a satisfying assignment for the above formula?
- (B) Describe a linear time algorithm that computes a satisfying assignment if it exists (hint: think about brooms!).

### 3.2. REDUCTIONS.

Show that the following problems can be reduced to the standard shortest path problems. No proof required.

- (A) Given directed graph  $G = (V, E)$  and two disjoint sets of nodes  $S, T$ . Find the shortest path from some node in  $S$  to some node in  $T$ .
- (B)  $G$  is a directed graph and nodes and edges have non-negative lengths. Find  $s$ - $t$  shortest path where the length of a path is equal to the sum of the lengths of the nodes and edges on the path.
- (C) Given a directed graph  $G$  with node lengths (no edge lengths), is there a negative length cycle? Here the length of a cycle is the sum of the lengths of nodes on the cycle.
- (D)  $G$  is a DAG and each node has a non-negative length. Given two nodes  $s, t$  in  $G$ , find the  $s$ - $t$  longest simple path in linear time.

### 3.3. QUICK FIX.

Your “friend” suggests that the easiest algorithm for finding shortest paths in a directed graph with negative-weighted edges is to make all the weights positive by adding a sufficiently large constant to each weight and then running Dijkstra’s algorithm. Give an example that you can show your friend to prove that his or her method is incorrect.

### 3.4. ALMOST POSITIVE.

We are given a directed graph  $G = (V, E)$  with potentially negative edge lengths. Your friend ran Dijkstra’s algorithm and came up with a shortest path tree  $T$  for distances from a node  $s$ . You realize that Dijkstra’s algorithm may not output distances correctly when a graph has negative edge lengths. However, before you run the more expensive Bellman-Ford algorithm, you wish to check whether  $T$  is a correct shortest path tree or not. Describe an

$O(m + n)$  time algorithm to do this check. Don't forget to prove that your algorithm is correct!

### 3.5. LIMITED SHORTEST PATHS.

We are given a directed graph in which the shortest path between any two vertices  $u$  and  $v$  is guaranteed to have at most  $k$  edges. Give an algorithm that finds the shortest path between two vertices  $u$  and  $v$  in  $O(k(n + m))$  time. Remember, edges can have negative weights.

### 3.6. AVERAGE CYCLE.

You are given a directed weighted graph  $G$  (the weights are positive), and a number  $x$ . Design an algorithm that decides if  $G$  has a cycle with average cost strictly smaller than  $x$ . The average cost of a cycle is the total weight of its edges divided by the number of edges. How fast is your algorithm?

### 3.7. BEST EDGE TO ADD, FAST.

Suppose you are given a directed graph  $G = (V, E)$  with non-negative edge lengths;  $\ell(e)$  is the length of  $e \in E$ . You are interested in the shortest path distance between two given locations/nodes  $s$  and  $t$ . It has been noticed that the existing shortest path distance between  $s$  and  $t$  in  $G$  is not satisfactory and there is a proposal to add exactly one edge to the graph to improve the situation. The candidate edges from which one has to be chosen is given by  $E' = \{e_1, e_2, \dots, e_k\}$  and you can assume that  $E \cap E' = \emptyset$ . The length of the  $e_i$  is  $\alpha_i \geq 0$ . Your goal is figure out which of these  $k$  edges will result in the most reduction in the shortest path distance from  $s$  to  $t$ . Describe an algorithm for this problem that runs in time  $O(n \log n + m + k)$  where  $m = |E|$  and  $n = |V|$ . Note that one can easily solve this problem in  $O(k(m + n) \log n)$  by running Dijkstra's algorithm  $k$  times, one for each  $G_i$  where  $G_i$  is the graph obtained by adding  $e_i$  to  $G$ .