

Concentration of Random Variables – Chernoff’s Inequality

15.1. Concentration of mass

15.1.1. Example: Binomial distribution

Consider the binomial distribution $\text{Bin}(n, 1/2)$ for various values of n as depicted in [Figure 15.1](#) – here we think about the value of the variable as the number of heads in flipping a fair coin n times. Clearly, as the value of n increases the probability of getting a number of heads that is significantly smaller or larger than $n/2$ is tiny. Here we are interested in quantifying exactly how far we can deviate from this expected value. Specifically, if $X \sim \text{Bin}(n, 1/2)$, then we would be interested in bound-




Figure 15.1: The binomial distribution for different values of n . It pretty quickly concentrates around its expectation.

$X_i \in \{-1, +1\}$ $\mathbb{P}[X_i = -1] = \mathbb{P}[X_i = 1] = 1/2$	
$\mathbb{P}[Y \geq \Delta] \leq \exp\left(-\frac{\Delta^2}{2n}\right)$	Theorem 15.2.1
$\mathbb{P}[Y \leq -\Delta] \leq \exp\left(-\frac{\Delta^2}{2n}\right)$	Theorem 15.2.1
$\mathbb{P}[Y - n/2 \geq \Delta] \leq 2 \exp\left(-\frac{2\Delta^2}{n}\right)$	Corollary 15.2.2

$X_i \in \{0, 1\}$ $\mathbb{P}[X_i = 1] = p_i, \mathbb{P}[X_i = 0] = 1 - p_i$	$X_i \in \{0, 1\}$ $\mathbb{P}[X_i = 1] = p_i, \mathbb{P}[X_i = 0] = 1 - p_i$
$\delta \geq 0$ $P = \mathbb{P}[Y > (1 + \delta)\mu] < \left(\frac{e^\delta}{(1+\delta)^{1+\delta}}\right)^\mu$ Theorem 15.4.1	$\delta \in (0, 1)$ $\mathbb{P}[Y < (1 - \delta)\mu] < \left(\frac{e^{-\delta}}{(1-\delta)^{1-\delta}}\right)^\mu$ Theorem 15.4.3
$\delta \in (0, 1)$ $P < \exp(-\mu\delta^2/3)$ Lemma 15.4.5	$\mathbb{P}[Y < (1 - \delta)\mu] < \exp(-\mu\delta^2/2)$ Lemma 15.4.4
$\delta \in (0, 4)$ $P < \exp(-\mu\delta^2/4)$ Lemma 15.4.6	
$\delta \in (0, 6)$ $P < \exp(-\mu\delta^2/5)$ Lemma 15.4.7	
$\delta \geq 2e - 1$ $P < 2^{-\mu(1+\delta)}$ Lemma 15.4.8	$\Delta \geq 0$ $\mathbb{P}[Y - \mu \geq \Delta] \leq \exp(-2\Delta^2/n)$ $\mathbb{P}[Y - \mu \leq -\Delta] \leq \exp(-2\Delta^2/n)$ Corollary 15.5.5
$\delta \geq e^2$ $P < \exp(-\frac{1}{2}\mu\delta \ln \delta)$ Lemma 15.4.9	
$\varphi \in (0, 1]$ $\mathbb{P}\left[Y > (1 + \delta)\mu + \frac{3 \ln \varphi^{-1}}{\delta^2}\right] < \varphi.$ Lemma 15.4.10	$\tau \geq 1$ $\mathbb{P}[Y < \mu/\tau] < \exp\left(-\left[1 - \frac{1 + \ln \tau}{\tau}\right]\mu\right)$ Theorem 15.4.3

$X_i \in [0, 1]$	Arbitrary independent distributions
$\delta \in [0, 1]$ $\mathbb{P}[Y \geq (1 + \delta)\mu] \leq \exp(-\delta^2\mu/4)$	Theorem 15.5.6
$\mathbb{P}[Y \leq (1 - \delta)\mu] \leq \exp(-\delta^2\mu/2)$	
$\Delta \geq 0$ $\mathbb{P}[Y - \mu \geq \Delta] \leq \exp(-2\Delta^2/n)$	Corollary 15.5.5
$\mathbb{P}[Y - \mu \leq -\Delta] \leq \exp(-2\Delta^2/n).$	

$X_i \in [a_i, b_i]$	Arbitrary independent distributions
$\Delta \geq 0$ $\mathbb{P}[Y - \mu \geq \Delta] \leq 2 \exp\left(-\frac{2\Delta^2}{\sum_{i=1}^n (b_i - a_i)^2}\right)$	Theorem 15.6.3

Table 15.1: Summary of Chernoff type inequalities covered. Here we have n independent random variables $X_1, \dots, X_n$, $Y = \sum_i X_i$ and $\mu = \mathbb{E}[Y]$.




Figure 15.2: The “middle” of the binomial distribution for different values of n . It very quickly converges to the normal distribution (under appropriate rescaling and translation).

ing the probability $\mathbb{P}[X > n/2 + \Delta]$, where $\Delta = t\sigma_X = t\sqrt{n}/2$ (i.e., we are t standard deviations away from the expectation). For $t > 2$, this probability is roughly 2^{-t} , which is what we prove here.

More surprisingly, if you look only at the middle of the distribution, it looks the same after clipping away the uninteresting tails, see [Figure 15.2](#); that is, it looks more and more like the normal distribution. This is a universal phenomenon known as the *central limit theorem* – every sum of nicely behaved random variables behaves like the normal distribution. We unfortunately need a more precise quantification of this behavior, thus the following.

15.1.2. A restricted case of Chernoff inequality via games

15.1.2.1. Chernoff games

The game. Consider the game where a player starts with $Y_0 = 1$ dollars. At every round, the player can bet a certain amount x (fractions are fine). With probability half she loses her bet, and with probability half she gains an amount equal to her bet. The player is not allowed to go all in – because if she loses then the game is over. So it is natural to ask what her optimal

betting strategy is, such that at the end of the game she has as much money as possible.

Is the game pointless? So, let Y_{i-1} be the money the player has at the end of the $(i-1)$ th round, and she bets an amount $\psi_i \leq Y_{i-1}$ in the i th round. As such, at the end of the i th round, she has

$$Y_i = \begin{cases} Y_{i-1} - \psi_i & \text{LOSE: probability half} \\ Y_{i-1} + \psi_i & \text{WIN: probability half} \end{cases}$$

dollars. This game, in expectation, does not change the amount of money the player has. Indeed, we have

$$\mathbb{E}[Y_i | Y_{i-1}] = \frac{1}{2}(Y_{i-1} - \psi_i) + \frac{1}{2}(Y_{i-1} + \psi_i) = Y_{i-1}.$$

And as such, we have that $\mathbb{E}[Y_i] = \mathbb{E}[\mathbb{E}[Y_i | Y_{i-1}]] = \mathbb{E}[Y_{i-1}] = \dots = \mathbb{E}[Y_0] =$

1. In particular, $\mathbb{E}[Y_n] = 1$ – namely, on average, independent of the player’s strategy she is not going to make any money in this game (and she is allowed to change her bets after every round). Unless she is lucky^{15.1}...

What about a lucky player? The player believes she will get lucky and wants to develop a strategy to take advantage of it. Formally, she believes that she can win, say, at least $(1+\delta)/2$ fraction of her bets (instead of the predicted $1/2$) – for example, if the bets are in the stock market, she can improve her chances by doing more research on the companies she is investing in^{15.2}. Unfortunately, the player does not know which rounds she is going to be lucky in – so she still needs to be careful.

In search of a good strategy. Of course, there are many safe strategies the player can use, from not playing at all, to risking only a tiny fraction of her money at each round. In other words, our quest here is to find the best strategy that extracts the maximum benefit for the player out of her inherent luck.

^{15.1}“I would rather have a general who was lucky than one who was good.” – Napoleon Bonaparte.

^{15.2}“I am a great believer in luck, and I find the harder I work, the more I have of it.” – Thomas Jefferson.

Here, we restrict ourselves to a simple strategy – at every round, the player would bet β fraction of her money, where β is a parameter to be determined. Specifically, at the end of the i th round, the player would have

$$Y_i = \begin{cases} (1 - \beta)Y_{i-1} & \text{LOSE} \\ (1 + \beta)Y_{i-1} & \text{WIN.} \end{cases}$$

By our assumption, the player is going to win in at least $M = (1 + \delta)n/2$ rounds. Our purpose here is to figure out what the value of β should be so that the player gets as rich as possible^{15.1}. Now, if the player is successful in $\geq M$ rounds, out of the n rounds of the game, then the amount of money the player has, at the end of the game, is

$$\begin{aligned} Y_n &\geq (1 - \beta)^{n-M} (1 + \beta)^M \\ &= (1 - \beta)^{n/2 - (\delta/2)n} (1 + \beta)^{n/2 + (\delta/2)n} \\ &= \left((1 - \beta)(1 + \beta) \right)^{n/2 - (\delta/2)n} (1 + \beta)^{\delta n} \\ &= \left(1 - \beta^2 \right)^{n/2 - (\delta/2)n} (1 + \beta)^{\delta n} \\ &\geq \exp(-2\beta^2)^{n/2 - (\delta/2)n} \exp(\beta/2)^{\delta n} \\ &= \exp((- \beta^2 + \beta^2 \delta + \beta \delta/2)n). \end{aligned}$$

To maximize this quantity, we choose $\beta = \delta/4$ (there is a better choice, see [Lemma 15.1.6](#), but we use this value for the simplicity of exposition).

Thus, we have that $Y_n \geq \exp\left(\left(-\frac{\delta^2}{16} + \frac{\delta^3}{16} + \frac{\delta^2}{8}\right)n\right) \geq \exp\left(\frac{\delta^2}{16}n\right)$, proving the following.

Lemma 15.1.1. *Consider a Chernoff game with n rounds, starting with one dollar, where the player wins in $\geq (1 + \delta)n/2$ of the rounds. If the player bets $\delta/4$ fraction of her current money, at all rounds, then at the end of the game the player has at least $\exp(n\delta^2/16)$ dollars.*

Remark 15.1.2. Note, that [Lemma 15.1.1](#) holds if the player wins any $\geq (1 + \delta)n/2$ rounds. In particular, the statement does not require randomness by itself – for our application, however, it is more natural and interesting to think about the player's wins as being randomly distributed.

^{15.1}This optimal choice is known as the Kelly criterion, see [Remark 15.1.3](#).

Remark 15.1.3. Interestingly, the idea of choosing the best fraction to bet is an old and natural question arising in investments strategies, and the right fraction to use is known as the *Kelly criterion*, going back to Kelly’s work from 1956 [Kel56].

15.1.3. A weak form of Chernoff’s inequality

The above implies that if a player is lucky, then she is going to become filthy rich^{15.1}. Intuitively, this should be a pretty rare event – because if the player is rich, then (on average) many other people have to be poor. We are thus ready for the kill.

Theorem 15.1.4 (Chernoff’s inequality). *Let $X_1, \dots, X_n$ be n independent random variables, where $X_i = 0$ or $X_i = 1$ with equal probability. Then, for any $\delta \in (0, 1/2)$, we have that*

$$\mathbb{P}\left[\sum_i X_i \geq (1 + \delta)\frac{n}{2}\right] \leq \exp\left(-\frac{\delta^2}{16}n\right).$$

Proof: Imagine that we are playing the Chernoff game above, with $\beta = \delta/4$, starting with 1 dollar, and let Y_i be the amount of money at the end of the i th round. Here $X_i = 1$ indicates that the player won the i th round. We have, by Lemma 15.1.1 and Markov’s inequality, that

$$\begin{aligned} \mathbb{P}\left[\sum_i X_i \geq (1 + \delta)\frac{n}{2}\right] &\leq \mathbb{P}\left[Y_n \geq \exp\left(\frac{n\delta^2}{16}\right)\right] \leq \frac{\mathbb{E}[Y_n]}{\exp(n\delta^2/16)} \\ &= \frac{1}{\exp(n\delta^2/16)} = \exp\left(-\frac{\delta^2}{16}n\right). \end{aligned} \quad \color{red}{\text{🐉}}$$

This is crazy – so intuition maybe? If the player is $(1 + \delta)/2$ -lucky then she can make a lot of money; specifically, at least $f(\delta) = \exp(n\delta^2/16)$ dollars by the end of the game. Namely, beating the odds has significant monetary value, and this value grows quickly with δ . Since we are in a “zero-sum” game setting, this event should be very rare indeed. Under this interpretation, of course, the player needs to know in advance the value of δ – so imagine that she guesses it somehow in advance, or she plays the game in parallel with all the possible values of δ , and she settles on the instance that maximizes her profit.

^{15.1}Not that there is anything wrong with that – many of my friends are filthy.

Can one do better? No, not really. The Chernoff inequality is tight (this is a challenging homework exercise) up to the constant in the exponent. The best bound I know for this version of the inequality has $1/2$ instead of $1/16$ in the exponent. Note, however, that no real effort was taken to optimize the constants – this is not the purpose here.

15.1.3.1. Some low level boring calculations Above, we used the following well known facts.

Lemma 15.1.5. (A) **Markov's inequality.** For any non-negative random variable X and $t > 0$, we have

$$\mathbb{P}[X \geq t] \leq \mathbb{E}[X] / t.$$

(B) For any two random variables X and Y , we have that $\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X | Y]]$.

(C) For $x \in (0, 1)$, $1 + x \geq e^{x/2}$.

(D) For $x \in (0, 1/2)$, $1 - x \geq e^{-2x}$.

Lemma 15.1.6. The quantity $\exp((- \beta^2 + \beta^2 \delta + \beta \delta / 2)n)$ is maximal for $\beta = \frac{\delta}{4(1-\delta)}$.

Proof: We have to maximize $f(\beta) = -\beta^2 + \beta^2 \delta + \beta \delta / 2$ by choosing the correct value of β (as a function of δ , naturally). $f'(\beta) = -2\beta + 2\beta \delta + \delta / 2 = 0 \iff 2(\delta - 1)\beta = -\delta / 2 \iff \beta = \frac{\delta}{4(1-\delta)}$. 

15.2. A proof for the $-1/ + 1$ case

Theorem 15.2.1. Let $X_1, \dots, X_n$ be n independent random variables, such that $\mathbb{P}[X_i = 1] = \mathbb{P}[X_i = -1] = \frac{1}{2}$, for $i = 1, \dots, n$. Let $Y = \sum_{i=1}^n X_i$. Then, for any $\Delta > 0$, we have

$$\mathbb{P}[Y \geq \Delta] \leq \exp(-\Delta^2 / 2n).$$

Proof: Clearly, for an arbitrary $t > 0$, to be specified shortly, we have

$$\mathbb{P}[Y \geq \Delta] = \mathbb{P}[\exp(tY) \geq \exp(t\Delta)] \leq \frac{\mathbb{E}[\exp(tY)]}{\exp(t\Delta)},$$

the first part follows by the fact that $\exp(\cdot)$ preserves ordering (requiring $t > 0$), and the second part follows by the Markov inequality.

Observe that

$$\begin{aligned}\mathbb{E}[\exp(tX_i)] &= \frac{1}{2}e^t + \frac{1}{2}e^{-t} = \frac{e^t + e^{-t}}{2} \\ &= \frac{1}{2} \left(1 + \frac{t}{1!} + \frac{t^2}{2!} + \frac{t^3}{3!} + \dots \right) \\ &\quad + \frac{1}{2} \left(1 - \frac{t}{1!} + \frac{t^2}{2!} - \frac{t^3}{3!} + \dots \right) \\ &= \left(1 + \frac{t^2}{2!} + \frac{t^4}{4!} + \dots + \frac{t^{2k}}{(2k)!} + \dots \right),\end{aligned}$$

by the Taylor expansion of $\exp(\cdot)$. Note, that $(2k)! \geq (k!)2^k$, and thus

$$\mathbb{E}[\exp(tX_i)] = \sum_{i=0}^{\infty} \frac{t^{2i}}{(2i)!} \leq \sum_{i=0}^{\infty} \frac{t^{2i}}{2^i(i!)} = \sum_{i=0}^{\infty} \frac{1}{i!} \left(\frac{t^2}{2} \right)^i = \exp(t^2/2),$$

again, by the Taylor expansion of $\exp(\cdot)$. Next, by the independence of the X_i s, we have

$$\begin{aligned}\mathbb{E}[\exp(tY)] &= \mathbb{E} \left[\exp \left(\sum_i tX_i \right) \right] = \mathbb{E} \left[\prod_i \exp(tX_i) \right] \\ &= \prod_{i=1}^n \mathbb{E}[\exp(tX_i)] \leq \prod_{i=1}^n e^{t^2/2} = e^{nt^2/2}.\end{aligned}$$

We have $\mathbb{P}[Y \geq \Delta] \leq \frac{\exp(nt^2/2)}{\exp(t\Delta)} = \exp(nt^2/2 - t\Delta)$.

Next, by minimizing the above quantity for t , we set $t = \Delta/n$. We conclude,

$$\mathbb{P}[Y \geq \Delta] \leq \exp \left(\frac{n}{2} \left(\frac{\Delta}{n} \right)^2 - \frac{\Delta}{n} \Delta \right) = \exp \left(-\frac{\Delta^2}{2n} \right). \quad \color{red}{\text{🔗}}$$

By the symmetry of Y , we get the following:

Corollary 15.2.2. *Let $X_1, \dots, X_n$ be n independent random variables, such that $\mathbb{P}[X_i = 1] = \mathbb{P}[X_i = -1] = \frac{1}{2}$, for $i = 1, \dots, n$. Let $Y = \sum_{i=1}^n X_i$. Then, for any $\Delta > 0$, we have $\mathbb{P}[|Y| \geq \Delta] \leq 2 \exp(-\Delta^2/2n)$.*

Corollary 15.2.3. *Let $X_1, \dots, X_n$ be n independent coin flips, such that $\mathbb{P}[X_i = 0] = \mathbb{P}[X_i = 1] = \frac{1}{2}$, for $i = 1, \dots, n$. Let $Y = \sum_{i=1}^n X_i$. Then, for any $\Delta > 0$, we have $\mathbb{P}[|Y - n/2| \geq \Delta] \leq 2 \exp(-2\Delta^2/n)$.*

Proof: Consider the random variables $Z_i = 2X_i - 1 \in \{-1, +1\}$. We have that

$$\begin{aligned} \mathbb{P}[|Y - n/2| \geq \Delta] &= \mathbb{P}[|2Y - n| \geq 2\Delta] = \mathbb{P}\left[\left|\sum_{i=1}^n (2X_i - 1)\right| \geq 2\Delta\right] \\ &= \mathbb{P}\left[\left|\sum_{i=1}^n Z_i\right| \geq 2\Delta\right] \leq 2 \exp(-2\Delta^2/n), \end{aligned}$$

by [Corollary 15.2.2](#) applied to the independent random variables $Z_1, \dots, Z_n$. 

Remark 15.2.4. Before going any further, it might be instrumental to understand what these inequalities imply. Consider the case where X_i is either zero or one with probability half. In this case $\mu = \mathbb{E}[Y] = n/2$. Set $\Delta = t\sqrt{n}$ ($\sqrt{\mu}$ is approximately the standard deviation of X if $p_i = 1/2$). We have by [Corollary 15.2.3](#):

$$\begin{aligned} \mathbb{P}\left[\left|Y - \frac{n}{2}\right| \geq \Delta\right] &\leq 2 \exp(-2\Delta^2/n) = 2 \exp(-2(t\sqrt{n})^2/n) \\ &= 2 \exp(-2t^2). \end{aligned}$$

Thus, the Chernoff inequality implies exponential decay (i.e., $\leq 2^{-t}$) with t standard deviations, instead of just polynomial (i.e., $\leq 1/t^2$) by Chebyshev's inequality.

15.3. To read or not to read?

From this point on, this note deteriorates into very tedious low-level calculus calculations. It's a repository of proofs that the bounds stated are correct. This is useful to read if you are interested in understanding the low-level details of how such bounds are proved, or enjoy such math (a lot of the math requires cleverness). I do not think, however, it provides much intuition for the inequalities themselves. Read from this point on for your own enjoyment, or suffering. You have been warned!

Another warning. It is very easy to misuse Chernoff's inequality (and its pals), and get brilliant short incorrect proofs. Make sure that you verify carefully the conditions required to deploy a certain version of the inequality, before using it.

15.4. The Chernoff Bound — General Case

Here we present the Chernoff bound in a more general setting.

Theorem 15.4.1. *Let $X_1, \dots, X_n$ be n independent variables, where*

$$\mathbb{P}[X_i = 1] = p_i \quad \text{and} \quad \mathbb{P}[X_i = 0] = q_i = 1 - p_i,$$

for all i . Let $X = \sum_{i=1}^n X_i$ and $\mu = \mathbb{E}[X] = \sum_i p_i$. For any $\delta > 0$, we have

$$\mathbb{P}[X > (1 + \delta)\mu] < \left(e^\delta / (1 + \delta)^{1+\delta} \right)^\mu.$$

Proof: We have $\mathbb{P}[X > (1 + \delta)\mu] = \mathbb{P}[e^{tX} > e^{t(1+\delta)\mu}]$. By the Markov inequality, we have:

$$\mathbb{P}[X > (1 + \delta)\mu] < \frac{\mathbb{E}[e^{tX}]}{e^{t(1+\delta)\mu}}$$

On the other hand,

$$\mathbb{E}[e^{tX}] = \mathbb{E}[e^{t(X_1 + X_2 + \dots + X_n)}] = \mathbb{E}[e^{tX_1}] \dots \mathbb{E}[e^{tX_n}].$$

Namely,

$$\begin{aligned} \mathbb{P}[X > (1 + \delta)\mu] &< \frac{\prod_{i=1}^n \mathbb{E}[e^{tX_i}]}{e^{t(1+\delta)\mu}} = \frac{\prod_{i=1}^n ((1 - p_i)e^0 + p_i e^t)}{e^{t(1+\delta)\mu}} \\ &= \frac{\prod_{i=1}^n (1 + p_i(e^t - 1))}{e^{t(1+\delta)\mu}}. \end{aligned}$$

Let $y = p_i(e^t - 1)$. We know that $1 + y < e^y$ (since $y > 0$). Thus,

$$\mathbb{P}[X > (1 + \delta)\mu] < \frac{\prod_{i=1}^n \exp(p_i(e^t - 1))}{e^{t(1+\delta)\mu}} = \frac{\exp(\sum_{i=1}^n p_i(e^t - 1))}{e^{t(1+\delta)\mu}}$$

$$\begin{aligned}
&= \frac{\exp((e^t - 1) \sum_{i=1}^n p_i)}{e^{t(1+\delta)\mu}} = \frac{\exp((e^t - 1)\mu)}{e^{t(1+\delta)\mu}} \\
&= \left(\frac{\exp(e^t - 1)}{e^{t(1+\delta)}} \right)^\mu = \left(\frac{\exp(\delta)}{(1 + \delta)^{(1+\delta)}} \right)^\mu,
\end{aligned}$$

if we set $t = \log(1 + \delta)$. 

15.4.1. The lower tail

We need the following low level lemma.

Lemma 15.4.2. *For $x \in [0, 1)$, we have $(1 - x)^{1-x} \geq \exp(-x + x^2/2)$.*

Proof: For $x \in [0, 1)$, we have, by the Taylor expansion, that $\ln(1 - x) = -\sum_{i=1}^{\infty} (x^i/i)$. As such, we have

$$\begin{aligned}
(1 - x) \ln(1 - x) &= -(1 - x) \sum_{i=1}^{\infty} \frac{x^i}{i} = -\sum_{i=1}^{\infty} \frac{x^i}{i} + \sum_{i=1}^{\infty} \frac{x^{i+1}}{i} \\
&= -x + \sum_{i=2}^{\infty} \left(\frac{x^i}{i-1} - \frac{x^i}{i} \right) = -x + \sum_{i=2}^{\infty} \frac{x^i}{i(i-1)}.
\end{aligned}$$

This implies that $(1 - x) \ln(1 - x) \geq -x + x^2/2$, which implies the claim by exponentiation. 

Theorem 15.4.3. *Let $X_1, \dots, X_n$ be n independent random variables, where $\mathbb{P}[X_i = 1] = p_i$ and $\mathbb{P}[X_i = 0] = q_i = 1 - p_i$, for all i . For $X = \sum_{i=1}^n X_i$, its expectation is $\mu = \mathbb{E}[X] = \sum_i p_i$. For any $\delta \in (0, 1)$, we have that*

$$\mathbb{P}[X < (1 - \delta)\mu] < \left[\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right]^\mu.$$

For any $\tau > 1$, we have that

$$\mathbb{P}[X < \mu/\tau] \leq \exp\left(-\left(1 - \frac{1 + \ln \tau}{\tau}\right)\mu\right).$$

Proof: We follow the same proof template seen already. For $t = -\ln(1 - \delta) > 0$, we have $\mathbb{E}[\exp(-tX_i)] = (1 - p_i)e^0 + p_ie^{-t} = 1 - p_i + p_i(1 - \delta) = 1 - p_i\delta \leq \exp(-p_i\delta)$. As such, we have

$$\begin{aligned}\mathbb{P}[X < (1 - \delta)\mu] &= \mathbb{P}[-X > -(1 - \delta)\mu] \\ &= \mathbb{P}[\exp(-tX) > \exp(-t(1 - \delta)\mu)] \\ &\leq \frac{\prod_{i=1}^n \mathbb{E}[\exp(-tX_i)]}{\exp(-t(1 - \delta)\mu)} \leq \frac{\exp(-\sum_{i=1}^n p_i\delta)}{\exp(-t(1 - \delta)\mu)} \\ &= \left[\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right]^\mu.\end{aligned}$$

For the last inequality, set $\delta = 1 - 1/\tau$, and observe that

$$\begin{aligned}\mathbb{P}[X < (1 - \delta)\mu] &\leq \left[\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right]^\mu = \left[\frac{\exp(-1 + 1/\tau)}{(1/\tau)^{1/\tau}} \right]^\mu \\ &= \exp\left(-\left(1 - \frac{1 + \ln \tau}{\tau}\right)\mu\right).\end{aligned}$$

Lemma 15.4.4. *Let $X_1, \dots, X_n \in \{0, 1\}$ be n independent random variables, with $p_i = \mathbb{P}[X_i = 1]$, for all i . Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For all $\delta \geq 0$, we have $\mathbb{P}[X < (1 - \delta)\mu] < \exp(-\mu\delta^2/2)$.*

Proof: This alternative simplified form of [Theorem 15.4.3](#), follows readily from [Lemma 15.4.2](#), since

$$\mathbb{P}[X < (1 - \delta)\mu] \leq \left[\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right]^\mu \leq \left[\frac{e^{-\delta}}{\text{Exp}(-\delta + \delta^2/2)} \right]^\mu \leq \text{Exp}(-\mu\delta^2/2).$$

15.4.2. A more convenient form of Chernoff's inequality

Lemma 15.4.5. *Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For $\delta \in (0, 1)$, we have*

$$\mathbb{P}[X > (1 + \delta)\mu] < \exp(-\mu\delta^2/3).$$

Proof: By **Theorem 15.4.1**, it is sufficient to prove, for $\delta \in [0, 1]$, that

$$\begin{aligned} \left(\frac{e^\delta}{(1+\delta)^{1+\delta}} \right)^\mu &\leq \exp\left(-\frac{\mu\delta^2}{c}\right) \\ \Leftrightarrow \mu(\delta - (1+\delta)\ln(1+\delta)) &\leq -\mu\delta^2/c \\ \Leftrightarrow f(\delta) = \delta^2/c + \delta - (1+\delta)\ln(1+\delta) &\leq 0. \end{aligned}$$

We have

$$f'(\delta) = 2\delta/c - \ln(1+\delta). \quad \text{and} \quad f''(\delta) = 2/c - \frac{1}{1+\delta}.$$

For $c = 3$, we have $f''(\delta) \leq 0$ for $\delta \in [0, 1/2]$, and $f''(\delta) \geq 0$ for $\delta \in [1/2, 1]$. Namely, $f'(\delta)$ achieves its maximum either at 0 or 1. As $f'(0) = 0$ and $f'(1) = 2/3 - \ln 2 \approx -0.02 < 0$, we conclude that $f'(\delta) \leq 0$. Namely, f is a monotonically decreasing function in $[0, 1]$, which implies that $f(\delta) \leq 0$, for all δ in this range, thus implying the claim. 

Lemma 15.4.6. *Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For $\delta \in (0, 4)$, we have*

$$\mathbb{P}[X > (1+\delta)\mu] < \exp(-\mu\delta^2/4).$$

Proof: **Lemma 15.4.5** implies a stronger bound, so we need to prove the claim only for $\delta \in (1, 4]$. Continuing as in the proof of **Lemma 15.4.5**, for case $c = 4$, we have to prove that

$$f(\delta) = \delta^2/4 + \delta - (1+\delta)\ln(1+\delta) \leq 0,$$

where $f''(\delta) = 1/2 - \frac{1}{1+\delta}$.

For $\delta > 1$, we have $f''(\delta) > 0$. Namely $f(\cdot)$ is convex for $\delta \geq 1$, and it achieves its maximum on the interval $[1, 4]$ on the endpoints. In particular, $f(1) \approx -0.13$, and $f(4) \approx -0.047$, which implies the claim. 

Lemma 15.4.7. *Let $X_1, \dots, X_n$ be n independent random variables, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For $\delta \in (0, 6)$, we have*

$$\mathbb{P}[X > (1+\delta)\mu] < \exp(-\mu\delta^2/5).$$

Proof: **Lemma 15.4.6** implies a stronger bound, so we need to prove the claim only for $\delta \in (4, 6]$. Continuing as in the proof of **Lemma 15.4.5**, for case $c = 5$, we have to prove that

$$f(\delta) = \delta^2/5 + \delta - (1 + \delta) \ln(1 + \delta) \leq 0,$$

where $f''(\delta) = 2/5 - \frac{1}{1+\delta}$. For $\delta \geq 4$, we have $f''(\delta) > 0$. Namely $f(\cdot)$ is convex for $\delta \geq 4$, and it achieves its maximum on the interval $[4, 6]$ on the endpoints. In particular, $f(4) \approx -0.84$, and $f(6) \approx -0.42$, which implies the claim. 

Lemma 15.4.8. *Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For $\delta > 2e - 1$, we have $\mathbb{P}[X > (1 + \delta)\mu] < 2^{-\mu(1+\delta)}$.*

Proof: By **Theorem 15.4.1**, we have

$$\left(\frac{e}{1 + \delta}\right)^{(1+\delta)\mu} \leq \left(\frac{e}{1 + 2e - 1}\right)^{(1+\delta)\mu} \leq 2^{-(1+\delta)\mu},$$

since $\delta > 2e - 1$. 

Lemma 15.4.9. *Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $X = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[X] = \sum_i p_i$. For $\delta > e^2$, we have $\mathbb{P}[X > (1 + \delta)\mu] < \exp\left(-\frac{\mu\delta \ln \delta}{2}\right)$.*

Proof: Observe that

$$\begin{aligned} \mathbb{P}[X > (1 + \delta)\mu] &< \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}}\right)^\mu \\ &= \exp\left(\mu\delta - \mu(1 + \delta) \ln(1 + \delta)\right). \end{aligned} \tag{15.1}$$

As such, we have

$$\begin{aligned} \mathbb{P}[X > (1 + \delta)\mu] &< \exp\left(-\mu(1 + \delta)(\ln(1 + \delta) - 1)\right) \\ &\leq \exp\left(-\mu\delta \ln \frac{1 + \delta}{e}\right) \end{aligned}$$

$$\leq \exp\left(-\frac{\mu\delta \ln \delta}{2}\right),$$

since for $x \geq e^2$ we have that $\frac{1+x}{e} \geq \sqrt{x} \iff \ln \frac{1+x}{e} \geq \frac{\ln x}{2}$. 

15.4.2.1. Bound when the expectation is small

Lemma 15.4.10. *Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $Y = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[Y] = \sum_i p_i$. For $\delta \in (0, 1]$, and $\varphi \in (0, 1]$, we have*

$$\mathbb{P}\left[Y > (1 + \delta)\mu + \frac{3 \ln \varphi^{-1}}{\delta^2}\right] < \varphi.$$

Proof: Let $\xi = \delta + \frac{3 \ln \varphi^{-1}}{\mu \delta^2}$. If $\xi \geq 2e - 1 \approx 4.43$, by [Lemma 15.4.8](#), we have

$$\alpha = \mathbb{P}\left[Y > (1 + \delta)\mu + \frac{3 \ln \varphi^{-1}}{\delta^2}\right] = \mathbb{P}[Y > (1 + \xi)\mu] \leq 2^{-\mu(1+\xi)} < \varphi,$$

since $\mu(1 + \xi) > \mu\xi = \mu\delta + \frac{3 \ln \varphi^{-1}}{\delta^2} \geq 3 \ln \varphi^{-1} > \log_2 \varphi^{-1}$ (which implies $2^{-\mu(1+\xi)} < \varphi$), as $\delta \in (0, 1]$.

If $\xi \leq 6$, then by [Lemma 15.4.7](#), we have

$$\alpha = \mathbb{P}[Y > (1 + \xi)\mu] \leq \exp(-\mu\xi^2/5) \leq \varphi,$$

since

$$\begin{aligned} -\frac{\mu}{5}\xi^2 &= -\frac{\mu}{5}\left(\delta + \frac{3 \ln \varphi^{-1}}{\mu \delta^2}\right)^2 \\ &\leq -\frac{\mu}{5}\left(2 \cdot \delta \cdot \frac{3 \ln \varphi^{-1}}{\mu \delta^2}\right) = -\frac{6}{5\delta} \ln \varphi^{-1} \leq -\ln \varphi^{-1} = \ln \varphi. \end{aligned} \quad \text{🐦}$$

Example 15.4.11. Let $X_1, \dots, X_n$ be n independent Bernoulli trials, where $\mathbb{P}[X_i = 1] = p_i$, and $\mathbb{P}[X_i = 0] = 1 - p_i$, for $i = 1, \dots, n$. Let $Y = \sum_{i=1}^n X_i$, and $\mu = \mathbb{E}[Y] = \sum_i p_i$. Assume that $\mu \leq 1/2$. Setting $\delta = 1$, we have, for $t > 6$, that

$$\mathbb{P}[Y > 1 + t] \leq \mathbb{P}\left[Y > (1 + \delta)\mu + \frac{3 \ln \exp(t/3)}{\delta^2}\right] \leq \exp(-t/3),$$

by [Lemma 15.4.10](#).

15.5. A special case of Hoeffding's inequality

In this section, we prove yet another version of Chernoff inequality, where each variable is randomly picked according to its own distribution in the range $[0, 1]$. We prove a more general version of this inequality in [Section 15.6](#), but the version presented here does not follow from this generalization.

Theorem 15.5.1. *Let $X_1, \dots, X_n \in [0, 1]$ be n independent random variables, let $X = \sum_{i=1}^n X_i$, and let $\mu = \mathbb{E}[X]$. We have that $\mathbb{P}[X - \mu \geq \eta] \leq \left(\frac{\mu}{\mu + \eta}\right)^{\mu + \eta} \left(\frac{n - \mu}{n - \mu - \eta}\right)^{n - \mu - \eta}$.*

Proof: Let $s \geq 1$ be some arbitrary parameter. By the standard arguments, we have

$$\gamma = \mathbb{P}[X \geq \mu + \eta] = \mathbb{P}[s^X \geq s^{\mu + \eta}] \leq \frac{\mathbb{E}[s^X]}{s^{\mu + \eta}} = s^{-\mu - \eta} \prod_{i=1}^n \mathbb{E}[s^{X_i}].$$

By calculations, see [Lemma 15.5.7](#) below, one can show that $\mathbb{E}[s^{X_i}] \leq 1 + (s - 1) \mathbb{E}[X_i]$. As such, by the AM-GM inequality^{15.1}, we have that

$$\begin{aligned} \prod_{i=1}^n \mathbb{E}[s^{X_i}] &\leq \prod_{i=1}^n \left(1 + (s - 1) \mathbb{E}[X_i]\right) \\ &\leq \left(\frac{1}{n} \sum_{i=1}^n \left(1 + (s - 1) \mathbb{E}[X_i]\right)\right)^n = \left(1 + (s - 1) \frac{\mu}{n}\right)^n. \end{aligned}$$

Setting $s = \frac{(\mu + \eta)(n - \mu)}{\mu(n - \mu - \eta)} = \frac{\mu n - \mu^2 + \eta n - \eta \mu}{\mu n - \mu^2 - \eta \mu}$ we have that

$$\begin{aligned} 1 + (s - 1) \frac{\mu}{n} &= 1 + \frac{\eta n}{\mu n - \mu^2 - \eta \mu} \cdot \frac{\mu}{n} = 1 + \frac{\eta}{n - \mu - \eta} \\ &= \frac{n - \mu}{n - \mu - \eta}. \end{aligned}$$

^{15.1}The inequality between arithmetic and geometric means:
 $(\sum_{i=1}^n x_i)/n \geq \sqrt[n]{x_1 \cdots x_n}$.

As such, we have that

$$\begin{aligned}\gamma &\leq s^{-\mu-\eta} \prod_{i=1}^n \mathbb{E}[s^{X_i}] = \left(\frac{\mu(n-\mu-\eta)}{(\mu+\eta)(n-\mu)} \right)^{\mu+\eta} \left(\frac{n-\mu}{n-\mu-\eta} \right)^n \\ &= \left(\frac{\mu}{(\mu+\eta)} \right)^{\mu+\eta} \left(\frac{n-\mu}{n-\mu-\eta} \right)^{n-\mu-\eta}.\end{aligned}$$

Remark 15.5.2. Setting $s = (\mu + \eta)/\mu$ in the proof of **Theorem 15.5.1**, we have

$$\mathbb{P}[X - \mu \geq \eta] \leq \left(\frac{\mu}{\mu+\eta} \right)^{\mu+\eta} \left(1 + \left(\frac{\mu+\eta}{\mu} - 1 \right) \frac{\mu}{n} \right)^n = \left(\frac{\mu}{\mu+\eta} \right)^{\mu+\eta} \left(1 + \frac{\eta}{n} \right)^n.$$

Corollary 15.5.3. Let $X_1, \dots, X_n \in [0, 1]$ be n independent random variables, let $\bar{X} = \sum_{i=1}^n X_i/n$, $p = \mathbb{E}[\bar{X}] = \mu/n$ and $q = 1 - p$. Then, we have that $\mathbb{P}[\bar{X} - p \geq t] \leq \exp(nf(t))$, for

$$f(t) = (p+t) \ln \frac{p}{p+t} + (q-t) \ln \frac{q}{q-t}. \quad (15.2)$$

Theorem 15.5.4. Let $X_1, \dots, X_n \in [0, 1]$ be n independent random variables, let $\bar{X} = (\sum_{i=1}^n X_i)/n$, and let $p = \mathbb{E}[\bar{X}]$. We have that

$$\begin{aligned}\mathbb{P}[\bar{X} - p \geq t] &\leq \exp(-2nt^2) \quad \text{and} \\ \mathbb{P}[\bar{X} - p \leq -t] &\leq \exp(-2nt^2).\end{aligned}$$

Proof: Let $p = \mu/n$, $q = 1 - p$, and let $f(t)$ be the function from **Eq. (15.2)**, for $t \in (-p, q)$. Now, we have that

$$\begin{aligned}f'(t) &= \ln \frac{p}{p+t} + (p+t) \frac{p+t}{p} \left(-\frac{p}{(p+t)^2} \right) \\ &\quad - \ln \frac{q}{q-t} - (q-t) \frac{q-t}{q} \frac{q}{(q-t)^2} \\ &= \ln \frac{p}{p+t} - \ln \frac{q}{q-t} = \ln \frac{p(q-t)}{q(p+t)}.\end{aligned}$$

As for the second derivative, we have

$$f''(t) = \frac{q \cancel{(p+t)}}{p(q-t)} \cdot \frac{p}{q} \cdot \frac{(p+t)(-1) - (q-t)}{(p+t)^2}$$

$$= \frac{-p-t-q+t}{(q-t)(p+t)} = -\frac{1}{(q-t)(p+t)} \leq -4.$$

Indeed, $t \in (-p, q)$ and the denominator $(q-t)(p+t)$ is maximized for $t = (q-p)/2$, and as such $(q-t)(p+t) \leq (2q - (q-p))(2p + (q-p))/4 = (p+q)^2/4 = 1/4$.

Now, $f(0) = 0$ and $f'(0) = 0$, and by Taylor's expansion, we have that $f(t) = f(0) + f'(0)t + \frac{f''(x)}{2}t^2 \leq -2t^2$, where x is between 0 and t .

The first bound now readily follows from plugging this bound into **Corollary 15.5.3**. The second bound follows by considering the random variables $Y_i = 1 - X_i$, for all i , and plugging this into the first bound. Indeed, for $\bar{Y} = 1 - \bar{X}$, we have that $q = \mathbb{E}[\bar{Y}]$, and then $\bar{X} - p \leq -t \iff t \leq p - \bar{X} \iff t \leq 1 - q - (1 - \bar{Y}) = \bar{Y} - q$. Thus, $\mathbb{P}[\bar{X} - p \leq -t] = \mathbb{P}[\bar{Y} - q \geq t] \leq \exp(-2nt^2)$. 

Corollary 15.5.5. *Let $X_1, \dots, X_n \in [0, 1]$ be n independent random variables, let $Y = \sum_{i=1}^n X_i$, and let $\mu = \mathbb{E}[Y]$. For any $\Delta > 0$, we have $\mathbb{P}[Y - \mu \geq \Delta] \leq \exp(-2\Delta^2/n)$ and $\mathbb{P}[Y - \mu \leq -\Delta] \leq \exp(-2\Delta^2/n)$.*

Proof: For $\bar{X} = Y/n$, $p = \mu/n$, and $t = \Delta/n$, by **Theorem 15.5.4**, we have

$$\mathbb{P}[Y - \mu \geq \Delta] = \mathbb{P}[\bar{X} - p \geq t] \leq \exp(-2nt^2) = \exp(-2\Delta^2/n). \quad \text{🐦}$$

Theorem 15.5.6. *Let $X_1, \dots, X_n \in [0, 1]$ be n independent random variables, let $X = \sum_{i=1}^n X_i$, and let $\mu = \mathbb{E}[X]$. For any $\varepsilon \in [0, 1]$, we have that*

$$\begin{aligned} \mathbb{P}[X - \mu \geq \varepsilon\mu] &\leq \exp(-\varepsilon^2\mu/4) \quad \text{and} \\ \mathbb{P}[X - \mu \leq -\varepsilon\mu] &\leq \exp(-\varepsilon^2\mu/2). \end{aligned}$$

Proof: Let $p = \mu/n$, and let $g(x) = f(px)$, for $x \in [0, 1]$ and $xp < q$. As before, computing the derivative of g , we have

$$\begin{aligned} g'(x) &= pf'(xp) = p \ln \frac{p(q-xp)}{q(p+xp)} = p \ln \frac{q-xp}{q(1+x)} \\ &\leq p \ln \frac{1}{1+x} \leq -\frac{px}{2}, \end{aligned}$$

since $(q - xp)/q$ is maximized for $x = 0$, and $\ln \frac{1}{1+x} \leq -x/2$, for $x \in [0, 1]$, as can be easily verified^{15.1}. Now, $g(0) = f(0) = 0$, and by integration, we have that $g(x) = \int_{y=0}^x g'(y)dy \leq \int_{y=0}^x (-py/2)dy = -px^2/4$. Now, plugging into [Corollary 15.5.3](#), we get that the desired probability $\mathbb{P}[X - \mu \geq \varepsilon\mu]$ is

$$\begin{aligned} \mathbb{P}[\bar{X} - p \geq \varepsilon p] &\leq \exp(nf(\varepsilon p)) = \exp\left(ng(\varepsilon)\right) \\ &\leq \exp(-pn\varepsilon^2/4) = \exp(-\mu\varepsilon^2/4). \end{aligned}$$

As for the other inequality, set $h(x) = g(-x) = f(-xp)$. Then

$$\begin{aligned} h'(x) &= -pf'(-xp) = -p \ln \frac{p(q + xp)}{q(p - xp)} = p \ln \frac{q(1 - x)}{q + xp} \\ &= p \ln \frac{q - xq}{q + xp} = p \ln \left(1 - x \frac{p + q}{q + xp}\right) \\ &= p \ln \left(1 - x \frac{1}{q + xp}\right) \leq p \ln(1 - x) \leq -px, \end{aligned}$$

since $1 - x \leq e^{-x}$. By integration, as before, we conclude that $h(x) \leq -px^2/2$. Now, plugging into [Corollary 15.5.3](#), we get

$$\begin{aligned} \mathbb{P}[X - \mu \leq -\varepsilon\mu] &= \mathbb{P}[\bar{X} - p \leq -\varepsilon p] \leq \exp(nf(-\varepsilon p)) \leq \exp(nh(\varepsilon)) \\ &\leq \exp(-np\varepsilon^2/2) \leq \exp(-\mu\varepsilon^2/2). \end{aligned} \quad \color{blue}{\text{🔗}}$$

15.5.1. Some technical lemmas

Lemma 15.5.7. *Let $X \in [0, 1]$ be a random variable, and let $s \geq 1$. Then $\mathbb{E}[s^X] \leq 1 + (s - 1)\mathbb{E}[X]$.*

Proof: By convexity, since $s \geq 1$, the function $x \mapsto s^x$ is convex on $[0, 1]$, so $s^x \leq (1 - x)s^0 + xs^1 = 1 + (s - 1)x$ pointwise for all $x \in [0, 1]$. Taking expectations on both sides immediately yields $\mathbb{E}[s^X] \leq 1 + (s - 1)\mathbb{E}[X]$ for any random variable $X \in [0, 1]$. 🔗

^{15.1}Indeed, this is equivalent to $\frac{1}{1+x} \leq e^{-x/2} \iff e^{x/2} \leq 1 + x$, which readily holds for $x \in [0, 1]$.

15.6. Hoeffding's inequality

In this section, we prove a generalization of Chernoff's inequality. The proof is considerably more tedious, and it is included here for the sake of completeness.

Lemma 15.6.1. *Let X be a random variable. If $\mathbb{E}[X] = 0$ and $a \leq X \leq b$, then for any $s > 0$, we have $\mathbb{E}[e^{sX}] \leq \exp(s^2(b-a)^2/8)$.*

Proof: Let $a \leq x \leq b$ and observe that x can be written as a convex combination of a and b . In particular, we have

$$x = \lambda a + (1 - \lambda)b \quad \text{for} \quad \lambda = \frac{b - x}{b - a} \in [0, 1].$$

Since $s > 0$, the function $\exp(sx)$ is convex, and as such

$$e^{sx} \leq \frac{b - x}{b - a} e^{sa} + \frac{x - a}{b - a} e^{sb},$$

since we have that $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$ if $f(\cdot)$ is a convex function. Thus, for a random variable X , by linearity of expectation, we have

$$\begin{aligned} \mathbb{E}[e^{sX}] &\leq \mathbb{E}\left[\frac{b - X}{b - a} e^{sa} + \frac{X - a}{b - a} e^{sb}\right] = \frac{b - \mathbb{E}[X]}{b - a} e^{sa} + \frac{\mathbb{E}[X] - a}{b - a} e^{sb} \\ &= \frac{b}{b - a} e^{sa} - \frac{a}{b - a} e^{sb}, \end{aligned}$$

since $\mathbb{E}[X] = 0$.

Next, set $p = -\frac{a}{b - a}$ and observe that $1 - p = 1 + \frac{a}{b - a} = \frac{b}{b - a}$ and

$$-ps(b - a) = -\left(-\frac{a}{b - a}\right)s(b - a) = sa.$$

As such, we have

$$\begin{aligned} \mathbb{E}[e^{sX}] &\leq (1 - p)e^{sa} + pe^{sb} = (1 - p + pe^{s(b-a)})e^{sa} \\ &= (1 - p + pe^{s(b-a)})e^{-ps(b-a)} \\ &= \exp(-ps(b-a) + \ln(1 - p + pe^{s(b-a)})) \\ &= \exp(-pu + \ln(1 - p + pe^u)), \end{aligned}$$

for $u = s(b - a)$. Setting

$$\phi(u) = -pu + \ln(1 - p + pe^u),$$

we thus have $\mathbb{E}[e^{sX}] \leq \exp(\phi(u))$. To prove the claim, we will show that $\phi(u) \leq u^2/8 = s^2(b - a)^2/8$.

To see that, expand $\phi(u)$ about zero using Taylor's expansion. We have

$$\phi(u) = \phi(0) + u\phi'(0) + \frac{1}{2}u^2\phi''(\theta) \quad (15.3)$$

where $\theta \in [0, u]$, and notice that $\phi(0) = 0$. Furthermore, we have

$$\phi'(u) = -p + \frac{pe^u}{1 - p + pe^u},$$

and as such $\phi'(0) = -p + \frac{p}{1-p+p} = 0$. Now,

$$\phi''(u) = \frac{(1 - p + pe^u)pe^u - (pe^u)^2}{(1 - p + pe^u)^2} = \frac{(1 - p)pe^u}{(1 - p + pe^u)^2}.$$

For any $x, y \geq 0$, we have $(x+y)^2 \geq 4xy$ as this is equivalent to $(x-y)^2 \geq 0$. Setting $x = 1 - p$ and $y = pe^u$, we have that

$$\phi''(u) = \frac{(1 - p)pe^u}{(1 - p + pe^u)^2} \leq \frac{(1 - p)pe^u}{4(1 - p)pe^u} = \frac{1}{4}.$$

Plugging this into [Eq. \(15.3\)](#), we get that

$$\begin{aligned} \phi(u) &\leq \frac{1}{8}u^2 = \frac{1}{8}(s(b - a))^2 \quad \text{and} \\ \mathbb{E}[e^{sX}] &\leq \exp(\phi(u)) \leq \exp\left(\frac{1}{8}(s(b - a))^2\right), \end{aligned}$$

as claimed. 

Lemma 15.6.2. *Let X be a random variable. If $\mathbb{E}[X] = 0$ and $a \leq X \leq b$, then for any $s > 0$, we have*

$$\mathbb{P}[X > t] \leq \frac{\exp\left(\frac{s^2(b-a)^2}{8}\right)}{e^{st}}.$$

Proof: Using the same technique we used in proving Chernoff's inequality, we have that

$$\mathbb{P}[X > t] = \mathbb{P}[e^{sX} > e^{st}] \leq \frac{\mathbb{E}[e^{sX}]}{e^{st}} \leq \frac{\exp\left(\frac{s^2(b-a)^2}{8}\right)}{e^{st}}. \quad \text{🔗}$$

Theorem 15.6.3 (Hoeffding's inequality). *Let $X_1, \dots, X_n$ be independent random variables, where $X_i \in [a_i, b_i]$, for $i = 1, \dots, n$. Then, for the random variable $S = X_1 + \dots + X_n$ and any $\eta > 0$, we have*

$$\mathbb{P}\left[|S - \mathbb{E}[S]| \geq \eta\right] \leq 2 \exp\left(-\frac{2\eta^2}{\sum_{i=1}^n (b_i - a_i)^2}\right).$$

Proof: Let $Z_i = X_i - \mathbb{E}[X_i]$, for $i = 1, \dots, n$. Set $Z = \sum_{i=1}^n Z_i$, and observe that

$$\mathbb{P}[Z \geq \eta] = \mathbb{P}[e^{sZ} \geq e^{s\eta}] \leq \frac{\mathbb{E}[\exp(sZ)]}{\exp(s\eta)},$$

by Markov's inequality. Arguing as in the proof of Chernoff's inequality, we have

$$\begin{aligned} \mathbb{E}[\exp(sZ)] &= \mathbb{E}\left[\prod_{i=1}^n \exp(sZ_i)\right] = \prod_{i=1}^n \mathbb{E}[\exp(sZ_i)] \\ &\leq \prod_{i=1}^n \exp\left(\frac{s^2(b_i - a_i)^2}{8}\right), \end{aligned}$$

since the Z_i s are independent and by **Lemma 15.6.1**. This implies that

$$\begin{aligned} \mathbb{P}[Z \geq \eta] &\leq \exp(-s\eta) \prod_{i=1}^n e^{s^2(b_i - a_i)^2/8} \\ &= \exp\left(\frac{s^2}{8} \sum_{i=1}^n (b_i - a_i)^2 - s\eta\right). \end{aligned}$$

The upper bound is minimized for $s = 4\eta/(\sum_i (b_i - a_i)^2)$, implying

$$\mathbb{P}[Z \geq \eta] \leq \exp\left(-\frac{2\eta^2}{\sum (b_i - a_i)^2}\right).$$

The claim now follows by the symmetry of the upper bound (i.e., apply the same proof to $-Z$). 🔗

15.7. Bibliographical notes

The general technique of bounding tail probabilities of sums of independent random variables by applying Markov's inequality to the moment-generating function was introduced by Bernstein [Ber24]. Chernoff [Che52] proved the bound in the context of the asymptotic efficiency of hypothesis tests, noting that the bound itself was communicated to him by Herman Rubin. The convenient multiplicative forms for independent Poisson trials (Theorem 15.4.1 and Lemma 15.4.5) were introduced and popularized in the computer science literature by Angluin and Valiant [AV79]. Hoeffding [Hoe63] proved the general bound for bounded independent random variables (Theorem 15.6.3) as well as the stronger form for variables in $[0, 1]$ (Theorem 15.5.1).

Some of the exposition here follows more or less the exposition in [MR95]. Exercise 15.8.1 (without the hint) is from [Mat99]. McDiarmid [McD89] provides a survey of Chernoff type inequalities; Theorem 15.5.6 and Section 15.5 are taken from there (our proof has somewhat weaker constants).

A more general treatment of such inequalities and tools is provided by Dubhashi and Panconesi [DP09].

15.8. Exercises

Exercise 15.8.1 (The Chernoff inequality is tight.). Let $S = \sum_{i=1}^n S_i$ be a sum of n independent random variables each attaining values $+1$ and -1 with equal probability. Let $P(n, \Delta) = \mathbb{P}[S > \Delta]$. Prove that for $\Delta \leq n/C$,

$$P(n, \Delta) \geq \frac{1}{C} \exp\left(-\frac{\Delta^2}{Cn}\right),$$

where C is a suitable constant. That is, the well-known Chernoff bound $P(n, \Delta) \leq \exp(-\Delta^2/(2n))$ is close to the truth.

Exercise 15.8.2 (The Chernoff inequality is tight by direct calculations.).

For this question use only basic argumentation – do not use Stirling's formula, Chernoff inequality or any similar “heavy” machinery.

(A) Prove that
$$\sum_{i=0}^{n-k} \binom{2n}{i} \leq \frac{n}{4k^2} 2^{2n}.$$

Hint: Consider flipping a coin $2n$ times. Write down explicitly the probability of this coin to have at most $n - k$ heads, and use Chebyshev inequality.

- (B) Using (A), prove that $\binom{2n}{n} \geq 2^{2n}/4\sqrt{n}$ (which is a pretty good estimate).
- (C) Prove that $\binom{2n}{n+i+1} = \left(1 - \frac{2i+1}{n+i+1}\right) \binom{2n}{n+i}$.
- (D) Prove that $\binom{2n}{n+i} \leq \exp\left(\frac{-i(i-1)}{2n}\right) \binom{2n}{n}$.
- (E) Prove that $\binom{2n}{n+i} \geq \exp\left(-\frac{8i^2}{n}\right) \binom{2n}{n}$.
- (F) Using the above, prove that $\binom{2n}{n} \leq c \frac{2^{2n}}{\sqrt{n}}$ for some constant c (I got $c = 0.824\dots$ but any reasonable constant will do).
- (G) Using the above, prove that

$$\sum_{i=t\sqrt{n}+1}^{(t+1)\sqrt{n}} \binom{2n}{n-i} \leq c 2^{2n} \exp(-t^2/2).$$

In particular, conclude that when flipping a fair coin $2n$ times, the probability to get less than $n - t\sqrt{n}$ heads (for t an integer) is smaller than $c' \exp(-t^2/2)$, for some constant c' .

- (H) Let X be the number of heads in $2n$ coin flips. Prove that for any $\delta > 0$ sufficiently small, it holds that

$$\mathbb{P}[X < (1 - \delta)n] \geq \exp(-c''\delta^2n),$$

where c'' is some constant. Namely, the Chernoff inequality is tight in the worst case.

Exercise 15.8.3 (Tail inequality for geometric variables). Let $X_1, \dots, X_m$ be m independent random variables with geometric distribution with probability p (i.e., $\mathbb{P}[X_i = j] = (1-p)^{j-1}p$). Let $Y = \sum_i X_i$, and let $\mu = \mathbb{E}[Y] = m/p$. For any $\delta \in (0, 1]$, prove that $\mathbb{P}[Y \geq (1 + \delta)\mu] \leq \exp(-m\delta^2/8)$.

References

- [AV79] D. Angluin and L. G. Valiant. **Fast probabilistic algorithms for Hamiltonian circuits and matchings**. *Journal of Computer and System Sciences*, 18(2): 155–193, 1979.
- [Ber24] S. N. Bernstein. **On a modification of Chebyshev’s inequality and of the error formula of Laplace**. *Ann. Sci. Inst. Sav. Ukraine, Sect. Math.*, 1(4): 38–49, 1924.
- [Che52] H. Chernoff. **A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations**. *The Annals of Mathematical Statistics*, 23(4): 493–507, 1952.
- [DP09] D. P. Dubhashi and A. Panconesi. *Concentration of Measure for the Analysis of Randomized Algorithms*. Cambridge University Press, 2009.
- [Hoe63] W. Hoeffding. **Probability inequalities for sums of bounded random variables**. *Journal of the American Statistical Association*, 58(301): 13–30, 1963.
- [Kel56] J. L. Kelly. **A new interpretation of information rate**. *Bell Sys. Tech. J.*, 35(4): 917–926, 1956.
- [Mat99] J. Matoušek. *Geometric Discrepancy*. Vol. 18. Algorithms and Combinatorics. Springer, 1999.
- [McD89] C. McDiarmid. **On the method of bounded differences**. *Surveys in Combinatorics*. Ed. by J. Siemons. Cambridge University Press, 1989, pp. 148–188.
- [MR95] R. Motwani and P. Raghavan. *Randomized Algorithms*. Vol. 26. Cambridge, UK: Cambridge University Press, 1995, pp. 48–50.